

On the Existence of Multiple Positive Solutions for a Semilinear Problem in Exterior Domains *

Yinbin Deng

Department of Mathematics, Huazhong Normal University,
Wuhan, 430079, P.R.CHINA

Yi Li

Department of Mathematics, University of Iowa
Iowa City, IA 52242, USA

Qingji Yang

Department of Mathematics, University of Rochester
Rochester, NY 14627, USA

July 12, 1999

Abstract

In this paper, we study the existence and nonexistence of multiple positive solutions for problem

$$\begin{cases} \Delta u + K(x)u^p = 0 & \text{in } \Omega. \\ u > 0 & \text{in } \Omega, \quad u \in H_{\text{loc}}^1(\Omega) \cap C(\bar{\Omega}). \\ u|_{\partial\Omega} = 0, & u \rightarrow \mu > 0 \quad \text{as } |x| \rightarrow \infty \end{cases}$$

where $\Omega = \mathbb{R}^N \setminus \omega$ is an exterior domain in \mathbb{R}^N , $\omega \subset \mathbb{R}^N$ is a bounded domain with smooth boundary and $N > 2$. $\mu \geq 0$, $p > 1$ are some given constants. $K(x)$ satisfies: $K(x) \in C_{\text{loc}}^\alpha(\Omega)$ and $\exists C, \epsilon, M > 0$ such that, $|K(x)| \leq C|x|^\epsilon$ for any $|x| \geq M$, with

*Research supported by the Natural Science Foundation of China

$l \leq -2 - \epsilon$. Some existence and nonexistence of multiple solutions have been discussed under different assumptions on K .

Key words and phrases: multiple solutions, critical exponents, elliptic equations.

Email: ybdeng@public.wh.hb.cn, yi-li@math.uiowa.edu, qyang@math.rhchester.edu

AMS Classification: 35J10 35J20 35J60 35J65

1 Introduction

In this paper, we study the existence of multiple solutions for problem

$$\begin{cases} \Delta u + K(x)u^p = 0 & \text{in } \Omega, \\ u > 0 & \text{in } \Omega, \quad u \in H_{\text{loc}}^1(\Omega) \cap C(\bar{\Omega}). \\ u|_{\partial\Omega} = 0, \end{cases} \quad (1.1)$$

with the boundary condition $u \rightarrow \mu > 0$ as $|x| \rightarrow \infty$, where $\Omega = \mathbb{R}^N \setminus \omega$ is an exterior domain in \mathbb{R}^N , $\omega \subset \mathbb{R}^N$ is a bounded domain with smooth boundary and $N > 2$. $p > 1$ is a given constants. $K(x)$ satisfies:

(H₁) $K(x) \in C_{\text{loc}}^\alpha(\Omega)$, $K \not\equiv 0$ and $\exists C, \epsilon, M > 0$ such that, $|K(x)| \leq C|x|^l$ for any $|x| \geq M$,

with $l \leq -2 - \epsilon$.

Such a problem occur in various branches of mathematical physis and Geometry. For $K(x) \equiv |x|^l$, $\Omega = \mathbb{R}^N$ equation (1.1) is known as Lane-Emden equation, sometimes it is also referred to as the Emden-Fowler equation in astrophysics, where u represents the density of a single star. When $p = \frac{N+2}{N-2}$, $\Omega = \mathbb{R}^N$ and $n \geq 3$, equation (1.1) is called the conformal scalar curvature equation in \mathbb{R}^N . Let g be the usual metric in \mathbb{R}^N , the problem of finding a metric g_1 which is conformal to g (i.e. $g_1 = u^{\frac{4}{N-2}}g$, for some positive function u with scalar curvature \tilde{K} is equivalent to that to find a positive solution of (1.1) with $K = \frac{N-2}{4(N-1)}\tilde{K}$. For a detail overview on (1.1), we refer readers to the papers [N2], [LN1], [Z] and the references therein.

Equations like (1.1) has been studied by many mathematicians ([B], [CZ], [CL1-2], [DL1-2], [DLZ], [DN1-2], [Es], [G], [GE], [JPY], [KL], [LY1-2], [Lio], [WW], [Y], [YY], [ZC]). Ni ([N1]), Kenig and Ni ([KN]) proved existence theorems for (1.1) under the condition (H_1) . It is shown in ([N1]) that if K is nonnegative with $K \geq Cr^l$ for some $l > (N-2)(p-1) - 2$ at infinity, or if K is nonpositive with $-K \geq Cr^l$ for some $l > -2$ at infinity then (1.1) possesses no positive solutions, where $C > 0$. Lin ([Lin]) proved the existence for (1.1) under the condition that $|K| \leq \frac{\phi(|x|)}{|x|^2}$ at infinity with $\int^\infty \frac{\phi(r)}{r} dr < \infty$. Lin in [Lin] also proved a nonexistence result when K is nonpositive with $-K \geq Cr^{-2}$ at infinity. Other nonexistence results are given in [BLY] and [LN1]. In case of that $|K| \leq Cr^{(N-2)(p-1)-2-\varepsilon}$ at infinity for some positive constants C and ε , the existence and asymptotics of positive solutions are studied by many authors, here we only mention the results of, for example, Ni, Yosutani [NY], [LN1], [LN2] and Li [L2]. In the fast decay case $|K| \leq Cr^l$, $l < -2$, Ni showed that (1.1) possesses infinitely many positive solutions which are bounded from below by positive constants (see [N1] and [LN1]). Li and Ni ([LN1]) showed that, for positive bounded solution of (1.1), the limit $u_\infty = \lim_{x \rightarrow \infty} u(x)$ always exists for any $\varepsilon > 0$, furthermore, if $u_\infty = 0$, then

$$u(x) \leq \begin{cases} C|x|^{2-N} & \text{if } p > \frac{N+l}{N-2}, \\ C_\varepsilon|x|^{\frac{(1-\varepsilon)(l+2)}{1-p}} & \text{if } p \leq \frac{N+l}{N-2}, \end{cases}$$

and if $u_\infty > 0$, then

$$|u - u_\infty| \leq \begin{cases} C|x|^{2-N} & \text{if } l < -N, \\ C|x|^{2-N} \log |x| & \text{if } l = -N, \\ C|x|^{2+l} & \text{if } -N < l < -2, \end{cases}$$

at ∞ . These results are refined in [LN2] and [L2].

Recently, Zhao (see [Z]) studied the following problem:

$$\begin{cases} \Delta u + K(x)f(u) = 0 & \text{in } \Omega \\ u > 0 & \text{in } \Omega, \quad u \in H_{\text{loc}}^1(\Omega) \cap C(\bar{\Omega}) \\ u|_{\partial\Omega} = 0, & u \rightarrow \mu > 0 \quad \text{as } |x| \rightarrow \infty. \end{cases} \quad (1.1)_\mu$$

The existence of one positive solution of problem $(1.1)_\mu$ when f is superlinear at 0 was obtained with some assumptions (Green-tight function) on $K(x)$ for small $\mu > 0$. A natural and interesting problem is that how many solutions can be obtained for a given $\mu > 0$. There seems to have been little progress in this direction. The purpose of this paper is to discuss the existence and nonexistence of multiple solutions for problem $(1.1)_\mu$ for a given $\mu > 0$. The main results of this paper can be included in the following theorems:

Theorem 1.1. *Suppose (H_1) . Let $h(x)$ be a positive harmonic function in Ω satisfying*

$$h(x) \Big|_{\partial\Omega} = 0, \quad \lim_{|x| \rightarrow \infty} h(x) = 1.$$

Then

- (i) *If $K(x) \leq 0$, then for any $\mu > 0$, there exists a unique solution u_μ of $(1.1)_\mu$. In addition, $u_\mu \leq \mu h$ on Ω and u_μ is increasing with respect to μ .*
- (ii) *If $K(x) \geq 0$, $\exists \mu^* \in (0, +\infty)$ and $\mu^* < +\infty$, such that for $\mu > \mu^*$ there does not exist a solution of $(1.1)_\mu$; and for $\mu \in (0, \mu^*)$, there exists a minimal solution u_μ of $(1.1)_\mu$. In addition, u_μ is increasing with respect to μ , $u_\mu \geq \mu h$ in Ω and, as $\mu \rightarrow \mu^*$, u_μ increase to u_{μ^*} the minimal solution of $(1.1)_{\mu^*}$, and u_{μ^*} is unique.*
- (iii) *If $K(x)$ change sign, we can find a $\mu^* \in (0, +\infty)$ such that problem $(1.1)_\mu$ possesses at least one solution for all $\mu \in (0, \mu^*)$.*

Theorem 1.2. *Suppose that $p = \frac{N+2}{N-2}$, (H_1) , $0 \leq K(x) \in L^1(\Omega)$, and*

(H_2) $K(x) > 0$ in a neighborhood V of some point $x_0 \in \Omega$ such that

$$K(x_0) = \sup_{x \in \Omega} K(x)$$

and $K(x) = K(x_0) + 0(|x - x_0|^2)$ near x_0 . Then problem $(1.1)_\mu$ possesses at least two solutions u_μ and U_μ with $u_\mu < U_\mu$ if $\mu \in (0, \mu^)$, where μ^* is given by Theorem 1.1.*

This paper is organized as follows: We first give some Lemmas in Section 2, which will be used in the proof of Theorem 1.1. Then the existence and nonexistence of minimal solution for problem $(1.1)_\mu$ is given in Section 3 by the standary barrier method. Finally, the existence of the second solution for $(1.1)_\mu$ is given in Section 4 by using the variational method.

2 Preliminaries

In this Section, we will prove some Lemmas which will be used in the proof of Theorem 1.1.

Lemma 2.1. *Let f be a locally Hölder continuous function on Ω with the following decay property*

$$|f(x)| \leq C|x|^\ell \quad \text{at } \infty \quad (2.1)$$

with $C > 0$, $\ell < -2$, and w be the Newtonian potential of f , i.e.

$$w(x) = \int_{\Omega} G(x, y)f(y)dy .$$

where $G(x, y)$ is the Green function for Ω corresponding to the Laplacian $-\Delta$. Then $w(x)$ is well-defined and at ∞ we have

$$|w(x)| \leq \begin{cases} C|x|^{2-N} & \text{if } \ell < -N \\ C|x|^{2-N} \ln |x| & \text{if } \ell = -N \\ C|x|^{2+\ell} & \text{if } -N < \ell < -2 \end{cases} \quad (2.2)$$

Proof. This Lemma may be proved by standard arguments. We include a proof here for the sake of completeness.

From the definition of Green function, we can easily deduce that

$$G(x, y) \leq \frac{C_N}{|x - y|^{N-2}}. \quad (2.3)$$

where $C_N = (N(N - 2)\omega_N)^{-1}$ and ω_N is the volume of the unit ball in \mathbb{R}^N . Using this fact and (2.1) we can find a constant $C > 0$ such that

$$|w(x)| \leq C \int_{\Omega} \frac{1}{|x - y|^{N-2}(1 + |y|^{-\ell})} dy . \quad (2.4)$$

Thus $w(x)$ is well-defined. Next we decompose the integral (2.4) as follows.

$$\begin{aligned} |w(x)| &\leq \left(\int_{|y-x| \leq \frac{|x|}{2}} + \int_{\frac{|x|}{2} \leq |y-x| \leq 2|x|} + \int_{2|x| \leq |y-x|} \right) \frac{C}{|y - x|^{N-2}(1 + |y|^{-\ell})} dy \\ &\equiv I_1 + I_2 + I_3 \end{aligned}$$

where I_1, I_2 and I_3 are defined by the last equality. Same as [LN2] we can conclude that

$$\begin{aligned} I_1 &\leq \frac{C}{|x|^{-\ell}} \int_0^{\frac{|x|}{2}} \frac{1}{r^{N-2}} r^{N-1} dr = C|x|^{2+\ell}, \\ I_2 &\leq C \int_{|x|/2}^{2|x|} \frac{1}{r^{N-2} r^{-\ell}} r^{N-1} dr = C|x|^{2+\ell}, \\ I_3 &\leq \begin{cases} C|x|^{2-N} & \text{if } N + \ell < 0, \\ C|x|^{2-N}(\ell n|x| + 1) & \text{if } N + \ell = 0, \\ C|x|^{2-N}(1 + |x|^{N+\ell}) & \text{if } N + \ell > 0, \end{cases} \end{aligned}$$

Now, it is easy to see that (2.2) holds. \square

Lemma 2.2. *Under the assumption of Lemma 2.1, suppose v is a solution of*

$$\begin{cases} -\Delta v = f(x) & \text{in } \Omega, \\ v|_{\partial\Omega} = 0 \quad \lim_{|x| \rightarrow \infty} v(x) = \mu. \end{cases} \quad (2.5)$$

Then

$$v = \mu h(x) + \int_{\Omega} G(x, y) f(y) dy \quad (2.6)$$

where $h(x)$ is the positive harmonic function in Ω satisfying

$$h(x) \Big|_{\partial\Omega} = 0, \quad \lim_{|x| \rightarrow \infty} h(x) = 1 \quad (2.7)$$

and $G(x, y)$ is the Green functions for Ω corresponding to Δ .

Proof. From [Z] and Lemma 2.1, we can deduce that $h(x)$ exists with $0 < h < 1$ in Ω and the integral in (2.6) is well-defined. Set $w(x) = \int_{\Omega} G(x, y) f(y) dy$. For an arbitrary but fixed point $z \in \Omega$, choose R large enough such that $R > |z|$ and $\omega \subset B_R(0)$. Now we define

$$\begin{aligned} w_1(x) &= \int_{\Omega \cap B_R(0)} G(x, y) f(y) dy, \\ w_2(x) &= \int_{\mathbb{R}^N \setminus B_R(0)} G(x, y) f(y) dy. \end{aligned}$$

Then it is standard that

$$\Delta w_1(z) + f(z) = 0 \quad \text{and} \quad \Delta w_2(z) = 0.$$

Since $w = w_1 + w_2$ we have

$$\Delta w + f = 0 \quad \text{in } \Omega. \quad (2.8)$$

By Lemma 2.1 and the property of Green functions we have

$$w \Big|_{\partial\Omega} = 0, \quad \lim_{|x| \rightarrow \infty} w(x) = 0. \quad (2.9)$$

Therefore

$$\begin{cases} \Delta(v - w) = 0, \\ (v - w)|_{\partial\Omega} = 0, \quad \lim_{|x| \rightarrow \infty} (v - w) = \mu. \end{cases}$$

By the uniqueness of the above problem, we have

$$v - w = \mu h(x).$$

This gives (2.6). □

Theorem 2.3. *Suppose (H_1) and let u be a bounded solution of $(1.1)_\mu$, then*

$$|u(x) - \mu h(x)| \leq \begin{cases} C|x|^{2-N} \text{ at } \infty & \text{if } \ell < -N \\ C|x|^{2-N} \ell n|x| \text{ at } \infty & \text{if } \ell = -N \\ C|x|^{2+\ell} \text{ at } \infty & \text{if } -N < \ell < -2 \end{cases}$$

where $h(x)$ is the unique solution of (2.7).

The proof of the above theorem can come directly from Lemma 2.1 and Lemma 2.2.

Lemma 2.4. *Suppose (H_1) with $l = -\frac{N+2}{2} - \epsilon$ for some $\epsilon > 0$, $K(x) \geq 0$, $K(x) \not\equiv 0$ and u_μ be the solution of $(1.1)_\mu$. Then*

$$u_\mu(x) - \mu h(x) \in \mathcal{D}_0^{1,2}(\Omega)$$

where $h(x)$ is the unique solution of (2.7) and $\mathcal{D}_0^{1,2}(\Omega)$ is a Sobolev's space defining as the completion of $C_0^\infty(\Omega)$ in the norm $\int_\Omega |\nabla u|^2 dx = \|u\|^2$.

Proof. From Theorem 2.3, (2.7), and $(1.1)_\mu$ we can easily conclude that

$$\begin{cases} \Delta(u_\mu - \mu h(x)) + K(x)u_\mu^p = 0 \\ (u_\mu - \mu h(x))|_{\partial\Omega} = 0, \quad \lim_{|x| \rightarrow \infty} (u_\mu - \mu h(x)) = 0 \end{cases}$$

and

$$-\int_{\Omega} \Delta(u_{\mu} - \mu h(x))(u_{\mu} - \mu h(x))dx = \int_{\Omega} |\nabla(u_{\mu} - \mu h(x))|^2 dx.$$

Thus

$$\begin{aligned} \int_{\Omega} |\nabla(u_{\mu} - \mu h(x))|^2 dx &= \int_{\Omega} K(x)u_{\mu}^p(u_{\mu} - \mu h(x))dx \\ &= \int_{\Omega \cap B_R} K(x)u_{\mu}^p(u_{\mu} - \mu h(x))dx + \int_{R^N \setminus B_R} K(x)u_{\mu}^p(u_{\mu} - \mu h(x))dx \\ &\leq C + C_1 \int_R^{\infty} r^l s(r)r^{N-1} dr \\ &\leq +\infty \end{aligned}$$

if $l < -\frac{N+2}{2}$. Here

$$s(r) = \begin{cases} |r|^{2-N} & \text{if } l < -N, \\ |r|^{2-N} \ln |r|, & \text{if } l = -N, \\ |r|^{2+l}, & \text{if } -N < l < -2. \end{cases}$$

Remark 2.1. The conclusion of Lemma 2.4 still remain true if we replace the assumption $l = -\frac{N+2}{2} - \epsilon$ by $S(|x|)K(x) \in L^1$ near ∞ .

□

3 Existence of minimal solution

In this Section, we will give a complete proof of Theorem 1.1 by the standary barrier method.

Lemma 3.1. *Suppose (H_1) and $K(x) \geq 0, K(x) \not\equiv 0$, then there exists a constant $0 < \mu^* < \infty$ such that problem $(1.1)_{\mu}$ possesses a minimal solution for all $\mu \in (0, \mu^*)$ and no solution for problem $(1.1)_{\mu}$ for $\mu > \mu^*$.*

Proof. First of all, we prove that problem $(1.1)_{\mu}$ has a minimal solution if μ is small enough.

In fact, let $\varphi(x) = h(x) + \int_{\Omega} G(x, y)K(y)dy$. From Lemma 2.2, $\varphi(x)$ is a solution of

$$\begin{cases} -\Delta\varphi = K(x) & \text{in } \Omega \\ \varphi|_{\partial\Omega} = 0, \lim_{|x| \rightarrow \infty} \varphi(x) = 1 \end{cases} \quad (3.1)$$

Denoting $\varphi_\mu(x) = \mu\varphi(x)$, we have $\varphi_\mu(x) \geq \mu h(x)$ because $K(x) \geq 0$ in Ω . Then

$$\begin{cases} -\Delta\varphi_\mu - K(x)\varphi_\mu^p = K(x)(\mu - (\mu\varphi)^p) \geq 0 \\ \varphi_\mu|_{\partial\Omega} = 0, \lim_{|x| \rightarrow \infty} \phi_\mu = \mu \end{cases}$$

if μ is small enough. So $\bar{u} = \mu\varphi$ is a supersolution of $(1.1)_\mu$ if μ is small enough. It is easy to check that $\underline{u} = \mu h(x)$ is a subsolution of $(1.1)_\mu$ for all $\mu > 0$ and all positive supersolution of $(1.1)_\mu$ must be larger than or equal to μh . The method of sub and supersolution yields our first claim.

Next, we set

$$\mu^* = \sup\{\mu > 0, \mid \text{problem } (1.1)_\mu \text{ possesses at least one solution}\} \quad (3.2)$$

so that $\mu^* > 0$. For any $\mu \in (0, \mu^*)$, from the definition of μ^* , we can find an $\bar{\mu} > \mu$ such that problem $(1.1)_{\bar{\mu}}$ possesses a solution $u_{\bar{\mu}}$ and hence $u_{\bar{\mu}}$ is a supersolution of $(1.1)_\mu$. It is easy to verify that $\underline{u}_\mu = \mu h$ is a subsolution of $(1.1)_\mu$ for all $\mu > 0$ and all positive supersolution of $(1.1)_\mu$ must be larger than or equal to μh . Using monotone iteration we can get the minimal solution u_μ for all $\mu \in (0, \mu^*)$.

Now, we are going to prove that $\mu^* < +\infty$. In fact, if u_μ solves $(1.1)_\mu$, since $u_\mu \geq \mu h$ we have

$$\begin{cases} -\Delta(u_\mu - \mu h(x)) = -\Delta u_\mu = K(x)(u_\mu)^{p-1} \geq K(x)(\mu h)^{p-1}(u_\mu - \mu h(x)) & \text{in } \Omega \\ (u_\mu - \mu h(x)) > 0 & \text{in } \Omega, \\ (u_\mu - \mu h(x)) \in \mathcal{D}_0^{1,2}(\Omega) \end{cases}$$

Thus the first eigenvalue of $-\Delta - K(x)(\mu h)^{p-1}$ on $\mathcal{D}^{1,2}(\Omega)$ is positive and this is impossible for μ large.

From the definition of μ^* we know that there is no solution for problem $(1.1)_\mu$ if $\mu > \mu^*$. \square

Lemma 3.2. *Suppose H_1) with $l = -\frac{N+2}{2} - \epsilon$ and $K(x) \geq 0$, $K(x) \not\equiv 0$. u_μ be the minimal solution of $(1.1)_\mu$ for $\mu \in (0, \mu^*)$. Then the minimizing problem*

$$\sigma_\mu = \inf \left\{ \int_\Omega |\nabla w|^2 dx \mid w \in \mathcal{D}_0^{1,2}(\Omega), \int_\Omega pK(x)u_\mu^{p-1}w^2 dx = 1 \right\} \quad (3.3)$$

can be attained by a function $\psi_\mu > 0$ which satisfies the equation

$$\begin{cases} -\Delta w = \sigma p K(x) u_\mu^{p-1} w & \text{in } \Omega \\ w \in \mathcal{D}_0^{1,2}(\Omega) \end{cases} \quad (3.4)_\mu$$

with $\sigma = \sigma_\mu$. Furthermore, $\sigma_\mu > 1$ for all $\mu \in (0, \mu^*)$.

Proof. We first prove that the functional $\int_\Omega p K u_\mu^{p-1} w^2 dx$ is weakly sequentially compact. In fact, let $\{w_n\}$ is a bounded sequence in $\mathcal{D}_0^{1,2}(\Omega)$ with weak limit $w \in \mathcal{D}_0^1(\Omega)$, the boundedness of K and u_μ in Ω and the use of Hölder inequality in a ball B_R for a large R , and $B'_R = \mathbb{R}^N \setminus B_R$ give

$$\begin{aligned} & \int_\Omega K u_\mu^{p-1} |w_n - w|^2 dx \\ & \leq C_1 \int_{B_R \cap \Omega} |w_n - w|^2 dx + C \left(\int_{B'_R} |w_n - w|^{\frac{2N}{N-2}} dx \right)^{\frac{N-2}{N}} \left(\int_{B'_R} K(x)^{\frac{N}{2}} dx \right)^{\frac{2}{N}} \end{aligned}$$

where C, C_1 are positive constants, independent of w_n, w . It follows from the compactness of the embedding $\mathcal{D}_0^{1,2}(\Omega \cap B_R) \hookrightarrow L^2(\Omega \cap B_R)$ and assumption (H_1) we have

$$\begin{aligned} \int_\Omega K u_\mu^{p-1} (w_n - w)^2 dx & \leq C_1 \int_{B_R \cap \Omega} |w_n - w|^2 dx + C \int_R^\infty r^{-(2+\epsilon) \cdot \frac{N}{2}} r^{N-1} dx \\ & \leq \frac{\epsilon_1}{2} + \frac{\epsilon_1}{2} = \epsilon_1 \end{aligned}$$

for any $\epsilon_1 > 0$ if R and n are large enough. This gives us that the functional $\int_\Omega p K u_\mu^{p-1} w_n^2 dx$ is weakly sequentially compact. Consequently standard minimization procedure shows that σ_μ is attained by a function $\psi_\mu \geq 0$, $\psi_\mu \in \mathcal{D}_0^{1,2}(\Omega)$, satisfying $(3.4)_\mu$ with $\sigma = \sigma_\mu$. By assumption (H_1) we deduce $\sigma_\mu p K(x) u_\mu^{p-1}(x) |x|^\delta \in L^q(\Omega)$ for some $\delta > 0$ and $q > \frac{N}{2}$. Therefore a result of Egnel [E] implies that ψ_μ is bounded in Ω and $\psi_\mu = 0(|x|^{2-N})$ as $|x| \rightarrow \infty$ and standard Hölder estimates then imply that $\psi_\mu \in C_{\text{loc}}^{1,\alpha}(\Omega)$ for all $0 < \alpha < 1$.

Next, we prove $\sigma_\mu > 1$. In fact, for $\mu < \bar{\mu}$, $\mu, \bar{\mu} \in (0, \mu^*)$ problem $(1.1)_\mu$ and $(1.1)_{\bar{\mu}}$ have a minimal solution u_μ and $u_{\bar{\mu}}$ respectively. Because $u_{\bar{\mu}}$ is a supersolution of $(1.1)_\mu$, we have

$u_\mu \leq u_{\bar{\mu}}$. Set $v_{\bar{\mu}} = u_{\bar{\mu}} - \bar{\mu}h$, $v_\mu = u_\mu - \mu h$. From Lemma 2.5 we have

$$\begin{cases} -\Delta v_{\bar{\mu}} = K(x)(v_{\bar{\mu}} + \bar{\mu}h)^p & v_{\bar{\mu}} > 0 \text{ in } \Omega \\ v_{\bar{\mu}} \Big|_{\partial\Omega} = 0, \lim_{|x| \rightarrow \infty} v_{\bar{\mu}}(x) = 0 & \text{and } v_{\bar{\mu}} \in \mathcal{D}_0^{1,2}(\Omega) \end{cases}$$

$$\begin{cases} -\Delta v_\mu = K(x)(v_\mu + \mu h)^p & v_\mu > 0 \text{ in } \Omega \\ v_\mu \Big|_{\partial\Omega} = 0, \lim_{|x| \rightarrow \infty} v_\mu(x) = 0 & \text{and } v_\mu \in \mathcal{D}_0^{1,2}(\Omega) \end{cases}$$

and

$$-\Delta(v_{\bar{\mu}} - v_\mu) = K(x)[(v_{\bar{\mu}} + \bar{\mu}h)^p - (v_\mu + \mu h)^p] = K(x)(u_{\bar{\mu}}^p - u_\mu^p) \geq 0.$$

Maximum principle gives us that

$$v_{\bar{\mu}} - v_\mu > 0 \quad \text{in } \Omega. \quad (3.5)$$

Furthermore,

$$\begin{cases} -\Delta(v_{\bar{\mu}} - v_\mu) = K(x)(u_{\bar{\mu}}^p - u_\mu^p) \geq K(x)pu_\mu^{p-1}(v_{\bar{\mu}} - v_\mu + (\bar{\mu} - \mu)h) \\ (v_{\bar{\mu}} - v_\mu) \in \mathcal{D}_0^{1,2}(\Omega). \end{cases} \quad (3.6)$$

On the other hand,

$$\begin{cases} -\Delta\psi_\mu = \sigma_\mu K(x)pu_\mu^{p-1}\psi_\mu & \psi_\mu \geq 0 \text{ in } \Omega \\ \psi_\mu \in \mathcal{D}_0^{1,2}(\Omega). \end{cases} \quad (3.7)$$

Multiplying (3.6) by ψ_μ and (3.7) by $w \equiv u_{\bar{\mu}} - u_\mu$ we deduce

$$\int_\Omega \nabla w \nabla \psi_\mu dx \geq \int_\Omega pK(x)u_\mu^{p-1}(w + (\bar{\mu} - \mu)h)\psi_\mu dx$$

and

$$\int_\Omega \nabla \psi_\mu \nabla w dx = \sigma_\mu p \int_\Omega K(x)u_\mu^{p-1}\psi_\mu w dx.$$

Thus

$$\begin{aligned} \sigma_\mu p \int_\Omega K(x)u_\mu^{p-1}\psi_\mu w dx &\geq \int_\Omega pK(x)u_\mu^{p-1}w\psi_\mu + p(\bar{\mu} - \mu) \int_\Omega K(x)u_\mu^{p-2}h\psi_\mu dx \\ &> \int_\Omega pK(x)u_\mu^{p-1}w\psi_\mu dx \end{aligned}$$

which gives $\sigma_\mu > 1$. □

Lemma 3.3. *Suppose (H_1) , $K(x) \geq 0$, $K(x) \not\equiv 0$ and $K(x) \in L^1(\Omega)$. Then there exists a constant $C > 0$ independent of μ such that*

$$\|u_\mu - \mu h\|_{\mathcal{D}_0^{1,2}(\Omega)} \leq C \quad \text{for all } \mu \in (0, \mu^*)$$

where u_μ is the minimal solution of $(1.1)_\mu$ and h is the unique solution of (2.7) .

Proof. Set $v_\mu = u_\mu - \mu h$. From Lemma 2.4 we have

$$\begin{cases} -\Delta v_\mu = K(x)(v_\mu + \mu h)^p, \\ v_\mu \in \mathcal{D}_0^{1,2}(\Omega) \end{cases}. \quad (3.8)$$

From Lemma 3.2 and (3.8) we deduce

$$\int_{\Omega} |\nabla v_\mu|^2 dx = \int_{\Omega} K(x)(v_\mu + \mu h)^p v_\mu dx \quad (3.9)$$

$$\int_{\Omega} |\nabla v_\mu|^2 dx \geq \sigma_\mu p \int_{\Omega} K(x)(v_\mu + \mu h)^{p-1} v_\mu^2 dx \quad (3.10)$$

and hence

$$\begin{aligned} \sigma_\mu p \int_{\Omega} K(x)(v_\mu + \mu h)^{p-1} v_\mu^2 dx &\leq \int_{\Omega} K(x)(v_\mu + \mu h)^p v_\mu dx \\ &\leq \int_{\Omega} K(x)(v_\mu + \mu h)^{p-1} v_\mu^2 dx + \int_{\Omega} K(x)(v_\mu + \mu h)^{p-1} \mu h v_\mu dx. \end{aligned}$$

So, for any $\epsilon > 0$,

$$\begin{aligned} (p-1) \int_{\Omega} K(x)(v_\mu + \mu h)^{p-1} v_\mu^2 dx &\leq \int_{\Omega} K(x) \mu h (v_\mu + \mu h)^{p-1} v_\mu dx \\ &\leq C \int_{\Omega} (K(x) v_\mu^p + K(x) v_\mu) dx \\ &\leq C \left(\int_{\Omega} K(x) dx \right)^{\frac{1}{p+1}} \left(\int_{\Omega} K v_\mu^{p+1} dx \right)^{\frac{p}{p+1}} \\ &\quad + C \left(\int_{\Omega} K(x) dx \right)^{\frac{p}{p+1}} \left(\int_{\Omega} K(x) v_\mu^{p+1} dx \right)^{\frac{1}{p+1}} \\ &\leq C_\epsilon \int_{\Omega} K(x) dx + \epsilon \int_{\Omega} K(x) v_\mu^{p+1} dx \end{aligned}$$

by Hölder's inequality and Young's inequality. Taking $\epsilon > 0$ small enough we deduce

$$\int_{\Omega} K(x) v_\mu^{p+1} dx \leq C \int_{\Omega} K(x) dx \leq C_1. \quad (3.11)$$

From (3.9), (3.10) we also have

$$\int_{\Omega} |\nabla v_{\mu}|^2 dx \leq \frac{1}{p} \int_{\Omega} |\nabla v_{\mu}|^2 dx + \int_{\Omega} K(x) \mu h (v_{\mu} + \mu h)^{p-1} v_{\mu} dx$$

and hence

$$\begin{aligned} \left(1 - \frac{1}{p}\right) \int_{\Omega} |\nabla v_{\mu}|^2 dx &\leq C |\mu^*|^p \|h\|_{\infty}^p \int_{\Omega} K(x) v_{\mu} dx + C \mu^* \|h\|_{\infty} \int_{\Omega} K(x) v_{\mu}^p dx \\ &\leq C \left(\int_{\Omega} K(x) dx \right)^{\frac{1}{p+1}} \left(\int_{\Omega} K(x) v_{\mu}^{p+1} dx \right)^{\frac{p}{p+1}} \\ &\quad + C \left(\int_{\Omega} K(x) dx \right)^{\frac{p}{p+1}} \left(\int_{\Omega} K(x) v_{\mu}^{p+1} dx \right)^{\frac{1}{p+1}} \\ &\leq C \end{aligned}$$

because of (3.11) and that $K(x) \in L^1(\Omega)$. □

Lemma 3.4. *Let $h(x)$ be the solution of (2.7) and suppose H_1), then for any $\mu > 0$, there exists a unique solution u_{μ} of (1.1) $_{\mu}$ if $K(x) \leq 0$. In addition, $u_{\mu} \leq \mu h$ on Ω and u_{μ} is increasing in μ .*

Proof. We remark that μh is a supersolution of (1.1) $_{\mu}$ which satisfies

$$\begin{cases} -\Delta(\mu h) - K(x)(\mu h)^p \geq -\mu \Delta h = 0 & \text{in } \Omega \\ \mu h|_{\partial\Omega} = 0, \lim_{|x| \rightarrow \infty} \mu h(x) = \mu & \\ \mu h > 0 & \text{in } \Omega \end{cases} \quad (3.12)$$

Next, let $\psi(x) = \int_{\Omega} G(x, y) |K(y)| dy$; from Lemma 2.2, $\psi(x)$ is the positive solution of

$$\begin{cases} -\Delta v = |K(x)| \\ v|_{\partial\Omega} = 0, v(x) \rightarrow 0 \text{ as } |x| \rightarrow \infty \end{cases} \quad (3.13)$$

we set $\underline{u} = (\mu h - \lambda \psi)^+$ for some $\lambda > 0$. We then have by standard results

$$-\Delta \underline{u} \leq -\lambda |K(x)|_{\{\underline{u} > 0\}} \leq K(x) \underline{u}^p \text{ on } \Omega$$

if λ is chosen such that

$$\underline{u}^p \leq (\mu h)^p \leq \lambda.$$

where $h(x)$ is the solution of (2.7). Thus \underline{u} is a nontrivial subsolution satisfies $\underline{u} \leq \mu h$ and the existence part is complete.

The various uniqueness and comparison results are deduced from the following claim.

Let $v, w \in H_{\text{loc}}^1(\Omega) \cap C_b(\Omega)$ satisfy

$$\begin{aligned} -\Delta v + |K(x)|v^p &\leq 0 \quad \text{in } \Omega \quad v \geq 0 \text{ in } \Omega \quad \lim_{|x| \rightarrow \infty} v \leq \mu, v|_{\partial\Omega} = 0, \\ -\Delta w + |K(x)|w^p &\geq 0 \quad \text{in } \Omega \quad w \geq 0 \text{ in } \Omega \quad \lim_{|x| \rightarrow \infty} w \geq \mu, w|_{\partial\Omega} = 0, \end{aligned}$$

then $v \leq w$ on Ω .

Indeed, for all $\epsilon > 0$, we may find R large enough such that

$$v \leq (1 + \epsilon)w \equiv w_\epsilon \quad \text{for } |x| \geq R$$

since we have on $B_R \cap \Omega$

$$\begin{aligned} &-\Delta(w_\epsilon - v) + p|K(x)|w_\epsilon^{p-1}(w_\epsilon - v) \\ &\geq -\Delta(w_\epsilon - v) + |K(x)|(w_\epsilon^p - v^p) \\ &= -\Delta w_\epsilon + |K(x)|w_\epsilon^p - (-\Delta v + |K(x)|v^p) \geq 0. \end{aligned}$$

Since the first eigenvalue of $-\Delta + p|K(x)|w_\epsilon^{p-1}$ is positive (on $H_0^1(\Omega \cap B_R)$) we deduce $w_\epsilon \geq v$ in Ω . Let $\epsilon \rightarrow 0$ we obtain our claim. Using the above claim we can easily deduce the uniqueness and that $u_\mu \leq \mu h$ for all $\mu > 0$ and $u_{\mu_1} \leq u_{\mu_2}$ if $\mu_1 \leq \mu_2$. \square

Lemma 3.5. *Suppose (H_1) , if $K(x)$ change sign, we can find a positive constant μ^* such that problem $(1.1)_\mu$ possesses at least one solution.*

Proof. Consider problem

$$\begin{cases} -\Delta v = K(x)(v + \mu h)^p, & v > 0 \quad \text{in } \Omega, \\ v|_{\partial\Omega} = 0, \lim_{|x| \rightarrow \infty} v(x) = 0. \end{cases} \quad (3.14)$$

From Lemma 3.1, we can find a positive constant μ^* such that problem

$$\begin{cases} -\Delta v = K^+(x)(v + \mu h)^p & \text{in } \Omega, \\ v|_{\partial\Omega} = 0, \lim_{|x| \rightarrow \infty} v = 0, v > 0 & \text{in } \Omega \end{cases}$$

possess a minimal solution \bar{v} for all $\mu \in (0, \mu^*)$. From Lemma 3.4, problem

$$\begin{cases} -\Delta v = -K^-(x)(v + \mu h)^p & \text{in } \Omega, \\ v|_{\partial\Omega} = 0, \lim_{|x| \rightarrow \infty} v = 0, v < 0 & \text{in } \Omega. \end{cases}$$

possesses a unique solution \underline{v} for all $\mu > 0$. Then \bar{v} is a supersolution of (3.14) $_{\mu}$ and \underline{v} is a subsolution of (3.14) $_{\mu}$. Furthermore $v = \bar{v} - \underline{v}$ satisfies

$$\begin{cases} -\Delta v = K^+(x)(\bar{v} + \mu h)^p + K^-(x)(\underline{v} + \mu h)^p \geq 0, \\ v|_{\partial\Omega} = 0, \lim_{|x| \rightarrow \infty} v = 0, \end{cases}$$

maximum principle implies that $v > 0$. The existence of solution for (3.14) $_{\mu}$ with $K(x)$ change sign come from the method of super-subsolution. Suppose v_{μ} be the solution of (3.14) $_{\mu}$, then $u_{\mu} = v_{\mu} + \mu h$ is a solution of (1.1) $_{\mu}$ with $0 < \underline{v} + \mu h < u_{\mu} < \bar{v} + \mu h$. \square

Theorem 3.6. *Suppose (H_1) . Let h be the solution of (2.10), then*

- (i) *If $K(x) \leq 0$, for any $\mu > 0$, there exists a unique solution u_{μ} of (1.1) $_{\mu}$. In addition, $u_{\mu} \leq \mu h$ on Ω and u_{μ} is increasing in μ .*
- (ii) *If $K(x) \geq 0$, $\exists \mu^* \in (0, \infty]$ and $\mu^* < +\infty$ if $K(x) \not\equiv 0$, such that for $\mu > \mu^*$ there does not exist a solution of (1.1) $_{\mu}$ and for $\mu \in (0, \mu^*)$, there exists a minimal solution u_{μ} of (1.1) $_{\mu}$. In addition, u_{μ} is increasing in μ , $u_{\mu} \geq \mu h$ in Ω . Finally, if*

$$K(x) \in L^1(\Omega), \tag{3.15}$$

then as $\mu \rightarrow \mu^$, u_{μ} increase to u_{μ^*} the minimal solution of (1.1) $_{\mu^*}$, and u_{μ^*} is unique.*

- (iii) *If $K(x)$ change sign, we can find a $\mu^* \in (0, +\infty)$ such that problem (1.1) $_{\mu}$ possesses at least one solution for all $\mu \in (0, \mu^*)$.*

Proof. From the above lemmas, we only have to prove that problem (1.1) $_{\mu^*}$ has a unique solution under the assumption (3.15). Denote the corresponding solution of (1.1) $_{\mu}$ by u_{μ} . Let $v_{\mu} = u_{\mu} - \mu h$. From assumption (3.15) and Lemma 3.3, we know $v_{\mu} \in \mathcal{D}_0^{1,2}(\Omega)$ and

$$\|v_{\mu}\|_{\mathcal{D}_0^{1,2}(\Omega)} \leq C < +\infty \text{ for all } \mu \in (0, \mu^*)$$

where C is a positive constant independent of μ . We claim that

$$\int_{\Omega} v_{\mu}^q dx \leq C < \infty \quad (3.16)$$

for all $q \geq \frac{2N}{N-2}$, where C is some positive constant independent of N . First of all, we consider $p \in (1, \frac{N+2}{N-2})$, the subcritical case. We adapt the argument due to Brezis and Kato [BK] to deduce the above claim. In fact, v_{μ} is a solution of

$$\begin{cases} -\Delta v_{\mu} = K(x)(v_{\mu} + \mu h)^p \\ v_{\mu} \in \mathcal{D}_0^{1,2}(\Omega) \quad v_{\mu} > 0 \end{cases} \quad \text{in } \Omega \quad (3.17)_{\mu}$$

Let $i > 1$, multiplying (3.17) $_{\mu}$ by v_{μ}^i and integrating by parts we obtain

$$4i(1+i)^{-2} \int_{\Omega} |\nabla v_{\mu}^{\frac{1}{2}(1+i)}|^2 dx = \int_{\Omega} K(x)(v_{\mu} + \mu h)^p v_{\mu}^i dx .$$

We refer by Hölder's and Young's inequalities that

$$\begin{aligned} \int_{\Omega} K(x)(v_{\mu} + \mu h)^p v_{\mu}^i dx &\leq C \int_{\Omega} K(x)(v_{\mu}^p + (\mu h)^p) v_{\mu}^i dx \\ &\leq C \int_{\Omega} K(x) v_{\mu}^{p+i} dx + C \int_{\Omega} K(x) v_{\mu}^i dx \\ &\leq C \int_{\Omega} K(x) v_{\mu}^{p+i} dx + C \left(\int_{\Omega} |K(x)| dx \right)^{\frac{p}{p+i}} \left(\int_{\Omega} K(x) v_{\mu}^{p+i} dx \right)^{\frac{i}{p+i}} \\ &\leq C \int_{\Omega} K(x) v_{\mu}^{p+i} dx + C \\ &\leq C \int_{\Omega} v_{\mu}^{p+i} dx + C . \end{aligned}$$

Thus

$$4i(1+i)^{-2} \int_{\Omega} |\nabla v_{\mu}^{\frac{1}{2}(1+i)}|^2 dx \leq C \int_{\Omega} v_{\mu}^{p+i} dx + C . \quad (3.18)$$

Let $\epsilon > 0$, be arbitrary, then for $i \geq \frac{2N}{N-2} - p > 1$, we have

$$t^{p+i} \leq \epsilon t^{i + \frac{N+2}{N-2}} + C_{\epsilon} t^{\frac{2N}{N-2}} . \quad (3.19)$$

for all $\epsilon > 0$ and $t \geq 0$, because $i + \frac{N+2}{N-2} > p + i \geq \frac{2N}{N-2}$. Applying the Sobolev's inequality

and (3.17)—(3.19) we have

$$\begin{aligned}
\left(\int_{\Omega} v_{\mu}^q dx\right)^{\frac{N-2}{N}} &= \left(\int_{\Omega} \left(v_{\mu}^{\frac{1}{2}(1+i)}\right)^{\frac{2N}{N-2}} dx\right)^{\frac{N-2}{N}} \\
&\leq C \int_{\Omega} v_{\mu}^{p+i} dx + C \\
&\leq C_{\epsilon} \int_{\Omega} v_{\mu}^{i+\frac{N+2}{N-2}} dx + C_{\epsilon} \int_{\Omega} v_{\mu}^{\frac{2N}{N-2}} dx + C \\
&= C_{\epsilon} \int_{\Omega} v_{\mu}^{\frac{(N-2)q}{N}} \cdot v_{\mu}^{\frac{4}{N-2}} dx + C \\
&\leq C_{\epsilon} \left(\int_{\Omega} v_{\mu}^q dx\right)^{\frac{N-2}{N}} \left(\int_{\Omega} v_{\mu}^{\frac{2N}{N-2}} dx\right)^{\frac{2}{N}} + C
\end{aligned}$$

with $q = \frac{N(1+i)}{N-2}$. From Lemma 3.3 and Sobolev inequality we deduce $\{v_{\mu}\}$ is bounded in $L^q(\Omega)$ for large $q > 1$ if we choose ϵ small enough.

Now, we are going to deal with the case when $p = \frac{N+2}{N-2}$. Our method is a combination of ideas found in papers of Brezis and Kato [BK] and Egnell [E]. For $j \geq 1$, define $\varphi_j(t) = t^j$, $t \geq 0$ and $\psi_j(t) = \int_0^t [\varphi'_j(s)]^2 ds = \frac{j^2}{2j-1} t^{2j-1}$. Let $\mu \in (0, \mu^*)$ and v_{μ} be the corresponding minimal solution of (3.17) $_{\mu}$. From Lemma 3.2 we have

$$\int_{\Omega} \nabla v_{\mu} \nabla v dx \geq p \int_{\Omega} K(x)(v_{\mu} + \mu h)^{p-1} v dx \quad (3.20)$$

for all $v \in \mathcal{D}_0^{1,2}(\Omega)$. By Theorem 2.3, Lemma 2.4 and Remark 2.1 we know $\varphi_j(v_{\mu}) \in \mathcal{D}_0^{1,2}(\Omega)$.

We may choose $v = \varphi_j(v_{\mu})$ in (3.20) to obtain

$$\int_{\Omega} |\varphi'_j(v_{\mu})|^2 |\nabla v_{\mu}|^2 dx \geq p \int_{\Omega} K(x)(v_{\mu} + \mu h)^{p-1} \varphi_j^2(v_{\mu}) dx. \quad (3.21)$$

Since v_{μ} is a solution of (3.17) $_{\mu}$ and $\psi_j(v_{\mu}) \in \mathcal{D}_0^{1,2}(\Omega)$, we also have

$$\int_{\Omega} \psi'_j(v_{\mu}) |\nabla v_{\mu}|^2 dx = \int_{\Omega} K(x)(v_{\mu} + \mu h)^p \psi_j(v_{\mu}) dx. \quad (3.22)$$

From (3.22), (3.21) we obtain

$$p \int_{\Omega} K(x)(v_{\mu} + \mu h)^{p-1} v_{\mu}^{2j} \leq \frac{j^2}{2j-1} \left[\int_{\Omega} K(x)(v_{\mu} + \mu h)^{p-1} v_{\mu}^{2j} + \int_{\Omega} K(x)(v_{\mu} + \mu h)^{p-1} \mu h v_{\mu}^{2j-1} \right] \quad (3.23)$$

since $\frac{j^2}{2j-1} \geq 1$ and is increasing in j , we may choose $j > 1$ sufficiently close to 1 such that

$\frac{j^2}{2j-1} < p$ for $j \leq j_0$. Set $\alpha(j, p) = p - \frac{j^2}{2j-1} > 0$. Then (3.23) gives

$$\begin{aligned} \alpha(j, p) \int_{\Omega} K(x) v_{\mu}^{p+2j-1} dx &\leq \alpha(j, p) \int_{\Omega} K(x) (v_{\mu} + \mu h)^{p-1} v_{\mu}^{2j} dx \\ &\leq \frac{j^2}{2j-1} \int_{\Omega} K(x) (v_{\mu} + \mu h)^{p-1} \mu h v_{\mu}^{2j-1} dx \\ &\leq \frac{Cj^2}{2j-1} \left[\int_{\Omega} K(x) v_{\mu}^{p+2j-2} \mu h dx + \int_{\Omega} K(x) (\mu h)^p v_{\mu}^{2j-1} dx \right] \\ &\leq C \left[\int_{\Omega} K(x) v_{\mu}^{p+2j-2} dx + \int_{\Omega} K(x) v_{\mu}^{2j-1} dx \right]. \end{aligned}$$

because $\mu < \mu^*$, $K(x) \leq C$. Since

$$\begin{aligned} \int_{\Omega} K(x) v_{\mu}^{p+2j-2} dx &\leq C \left[\int_{\Omega} K(x) dx \right]^{\frac{1}{p+2j-1}} \left[\int_{\Omega} K(x) v_{\mu}^{p+2j-1} dx \right]^{\frac{p+2j-2}{p+2j-1}} \\ &\leq C \int_{\Omega} K(x) dx + \frac{\delta}{2} \int_{\Omega} K(x) v_{\mu}^{p+2j-1} dx \end{aligned}$$

for all $\delta > 0$ and similarly,

$$\int_{\Omega} K(x) v_{\mu}^{2j-1} dx \leq C \int_{\Omega} K(x) dx + \frac{\delta}{2} \int_{\Omega} K(x) v_{\mu}^{p+2j-1} dx$$

for all $\delta > 0$, we can deduce

$$(\alpha(j, p) - \delta) \int_{\Omega} K(x) v_{\mu}^{p+2j-1} dx \leq C_{\delta} \int_{\Omega} K(x) dx .$$

From the assumption of $K(x) \in L^1(\Omega)$, we have

$$\int_{\Omega} K(x) v_{\mu}^{p+2j-1} dx \leq C_{\delta} \tag{3.23}^*$$

for $j \in (1, j_0]$ and $C > 0$ independent of $\mu \in (0, \mu^*)$ if we take δ small enough. This shows that (3.16) holds for all $q \in [\frac{2N}{N-2}, p + 2j_0 - 1]$. To establish (3.16) for all $q \geq \frac{2N}{N-2}$ we use ideas in Brezis and Kato [BK]. Set $q_0 = \frac{2N}{N-2}$, $\delta = p + 2j_0 - 1 - \frac{2N}{N-2} > 0$. Multiplication of (3.17) by $v_{\mu}^{q_0-1}$, integration by parts and simple application of Hölder's inequality and

Young's inequality yield

$$\begin{aligned}
(q_0 - 1)q_0^{-2} \int |\nabla v_\mu^{\frac{q_0}{2}}|^2 dx &= \int_\Omega K(x)(v_\mu + \mu h)^p v_\mu^{q_0-1} dx \\
&\leq C \int_\Omega K(x)v_\mu^{p+q_0-1} dx + C \int_\Omega K(x)v_\mu^{q_0-1} dx \\
&\leq C \int_\Omega K(x)v_\mu^{p+q_0-1} dx \\
&\quad + C \left(\int_\Omega K(x) dx \right)^{\frac{p}{p+q_0-1}} \left(\int_\Omega K(x)v_\mu^{p+q_0-1} dx \right)^{\frac{q_0-1}{p+q_0-1}} \\
&\leq C \int_\Omega K(x)v_\mu^{p+q_0-1} dx + C \int_\Omega K(x) dx
\end{aligned}$$

which gives us

$$(q_0 - 1)q_0^{-2} \int |\nabla v_\mu^{\frac{q_0}{2}}|^2 dx \leq C \int_\Omega K(x)v_\mu^{p+q_0-1} dx + C \int_\Omega K(x) dx \quad (3.24)$$

where C is a positive constant independent of μ .

For any given $\epsilon > 0$ we can find a positive constant C_ϵ such that

$$v_\mu^{p-1+q_0} \leq \epsilon v^{p-1+q_0+\frac{2\delta}{N}} + C_\epsilon v_\mu^{q_0}.$$

This can be easily verified by the fact that $q_0 < p - 1 + q_0 < p - 1 + q_0 + \frac{2\delta}{N}$. Therefore, it follows from Hölder inequality, Sobolev's inequality and (3.23)* with $j = j_0$ that

$$\begin{aligned}
\int_\Omega K(x)v_\mu^{p+q_0-1} dx &\leq \epsilon \int_\Omega K(x)v_\mu^{p-1+q_0+\frac{2\delta}{N}} dx + C_\epsilon C \\
&\leq \epsilon \left(\int_\Omega K(x)(v_\mu^{q_0})^{\frac{p+1}{2}} dx \right)^{\frac{2}{p+1}} \left(\int_\Omega K(x)v_\mu^{p+2j_0-1} dx \right)^{\frac{2}{N}} + C_\epsilon \\
&\leq \epsilon C \int_\Omega (v_\mu^{\frac{q_0}{2}})^{p+1} dx + C_\epsilon C \\
&\leq \epsilon C \int_\Omega |\nabla v_\mu^{\frac{q_0}{2}}|^2 dx + C_\epsilon C.
\end{aligned}$$

which gives us

$$\int_\Omega K(x)v_\mu^{p+q_0-1} dx \leq \epsilon C \int_\Omega |\nabla v_\mu^{\frac{q_0}{2}}|^2 dx + C_\epsilon C. \quad (3.25)$$

It follows from (3.24) and (3.25), with ϵ sufficiently small, that

$$\int_\Omega |\nabla v_\mu^{\frac{q_0}{2}}|^2 dx \leq C \quad (3.26)$$

for some constant C , independent of μ , and by Sobolev's inequality we have

$$\int_{\Omega} v_{\mu}^{\frac{q_0}{2} \cdot q_0} dx \leq C .$$

The desired inequality (3.16) then follows easily by iteration. Set $q_1 = \frac{q_0^2}{2}$ and $q_k = \frac{q_{k-1}^2}{2}$.

Denote $g_{\mu}(x) = K(x)(v_{\mu}(x) + \mu h(x))^p$. From the above proof we deduce $g_{\mu}(x) \in L^q(\Omega)$ for all $q \geq p+1$ and

$$\begin{aligned} \int_{\Omega} |g_{\mu}(x)|^q dx &\leq C \int_{\Omega} K(x)^q v_{\mu}(x)^{pq} dx + C \int_{\Omega} K(x)^q (\mu h(x))^{pq} dx \\ &\leq C |K(x)|_{\infty}^{q-1} \int_{\Omega} K(x) v_{\mu}^{pq} dx + C \mu^* |h(x)|_{\infty}^q \int_{\Omega} K(x)^q dx \\ &\leq C \end{aligned}$$

for all $\mu \in (0, \mu^*)$.

We employ a classical a priori estimate to obtain

$$\|v_{\mu}\|_{\infty, B_R(x) \cap \Omega} \leq C_R (\|v_{\mu}\|_{p+1, B_{2R}(x) \cap \Omega} + \|g_{\mu}\|_{q, B_{2R}(0) \cap \Omega})$$

for solution of $-\Delta v = g_{\mu}(x)$, where $B_R(x)$ is a ball of radius R and centre x , and C_R is a constant independent of μ and x . Hölder estimates in $B_R \cap \Omega$ then shows that

$$\|v_{\mu}\|_{C^{1,\alpha}(B_R \cap \Omega)} \leq C_R$$

for some constant C_R , independing of μ . A simple diagonalization argument and the Ascoli-Arzela theorem may be employed to show that for a subsequence $\mu_n \rightarrow \mu^*$, $v_{\mu_n}, |\nabla v_{\mu_n}|$ converge uniformly on each compact subset of Ω , to a function $v_{\mu^*} \in \mathcal{D}_0^{1,2}(\Omega)$. It follows that

$$\int_{\Omega} \nabla v \cdot \nabla v_{\mu^*} dx = \int_{\Omega} K(x)(v_{\mu^*} + \mu^* h)^p v dx$$

for all $v \in C_0^{\infty}(\Omega)$ and therefore v_{μ^*} is a nonnegative weak solution of (3.17) $_{\mu^*}$. Thus $u_{\mu^*} = v_{\mu^*} + \mu^* h$ is a solution of (1.1) $_{\mu^*}$.

Finally, we prove that u_{μ^*} is unique. In fact, from the definition we can easily deduce that $\sigma_{\mu^*} = 1$ by applying the implicit function theorem to the function $F : \mathcal{D}_0^{1,2}(\Omega) \hookrightarrow \mathcal{D}_0^{1,2}(\Omega)$ with

$$F(u) = -\Delta u - K(x)(u + \mu h)^p \quad u \in \mathcal{D}_0^{1,2}(\Omega) .$$

If there exists another solution $\bar{u}_{\mu^*} \geq u_{\mu^*}$ for problem $(1.1)_{\mu^*}$, set $\bar{v}_{\mu^*} = \bar{u}_{\mu^*} - \mu^*h$, $v_{\mu^*} = u_{\mu^*} - \mu^*h$. We have from (3.17) $_{\mu}$

$$\begin{aligned} -\Delta(\bar{v}_{\mu^*} - v_{\mu^*}) &= K(x)[(\bar{v}_{\mu^*} + \mu^*h)^p - (v_{\mu^*} + \mu^*h)^p] \\ &= K(x)[p(v_{\mu^*} + \mu^*h)^{p-1}(\bar{v}_{\mu^*} - v_{\mu^*}) \\ &\quad + p(p-1)(v_{\mu^*} + \theta(\bar{v}_{\mu^*} - v_{\mu^*}) + \mu^*h)^{p-2}(\bar{v}_{\mu^*} - v_{\mu^*})] \end{aligned}$$

for some $\theta(x) \in [0, 1]$. From Lemma 3.2 and the above equality, we deduce

$$\begin{aligned} \sigma(\mu^*) \int_{\Omega} p(v_{\mu^*} + \mu^*h)^{p-1}(\bar{v}_{\mu^*} - v_{\mu^*})\psi_{\mu^*} dx &= \int_{\Omega} \nabla\psi_{\mu^*}\nabla(\bar{v}_{\mu^*} - v_{\mu^*}) dx \\ &= \int_{\Omega} p(v_{\mu^*} + \mu^*h)^{p-1}(\bar{v}_{\mu^*} - v_{\mu^*})\psi_{\mu^*} dx \\ &\quad + \int_{\Omega} p(p-1)(v_{\mu^*} + \theta(\bar{v}_{\mu^*} - v_{\mu^*}) + \mu^*h)^{p-2}(\bar{v}_{\mu^*} - v_{\mu^*})^2\psi_{\mu^*} dx \\ \text{i.e. } (\sigma(\mu^*) - 1) \int_{\Omega} p(v_{\mu^*} + \mu^*h)^{p-1}(\bar{v}_{\mu^*} - v_{\mu^*})\psi_{\mu^*} dx & \\ &= \int_{\Omega} p(p-1)(v_{\mu^*} + \theta(\bar{v}_{\mu^*} - v_{\mu^*}) + \mu^*h)^{p-2}(\bar{v}_{\mu^*} - v_{\mu^*})^2\psi_{\mu^*} dx \end{aligned}$$

we can obtain that $\bar{v}_{\mu^*} \equiv v_{\mu^*}$ from $\sigma(\mu^*) = 1$. □

4 The existence of second solution

For $\mu \in (0, \mu^*)$, let u_{μ} be the first solution of $(1.1)_{\mu}$ and consider the problem

$$\begin{cases} -\Delta v = K(x)((v + u_{\mu})^p - u_{\mu}^p) & \text{in } \Omega \\ v \in \mathcal{D}_0^1(\Omega), \quad v > 0 & \text{in } \Omega. \end{cases} \quad (4.1)_{\mu}$$

It is clear that $U_{\mu} = v_{\mu} + u_{\mu}$ is a solution of $(1.1)_{\mu}$ if v_{μ} is a solution of $(4.1)_{\mu}$. Consider the energy functional J_{μ} defined by

$$J_{\mu}(v) = \int_{\Omega} \frac{1}{2} |\nabla v|^2 - K(x) \left[\frac{1}{p+1} (u_{\mu} + v^+)^{p+1} - \frac{1}{p+1} u_{\mu}^{p+1} - u_{\mu}^p v^+ \right] dx.$$

Standard procedure from the calculus of variations shows that J_{μ} is well defined in $\mathcal{D}_0^1(\Omega)$ with continuous Freehet derivative given by

$$J'_{\mu}(v)\varphi = \int_{\Omega} [\nabla v \nabla \varphi - K(x)((u_{\mu} + v^+)^p - u_{\mu}^p)]\varphi dx \quad \varphi \in \mathcal{D}_0^{1,2}(\Omega)$$

A critical point v of J_μ is a weak solution of the equation

$$-\Delta v = K(x)[(u_\mu + v^+)^p - u_\mu^p] \quad v \in \mathcal{D}_0^1(\Omega)$$

and if $v > 0$ in \mathbb{R}^N , then v is a solution of (4.1) $_\mu$.

The following Lemma comes from the fact that

$$\lim_{s \rightarrow 0} \frac{(u_\mu + s)^p - u_\mu^p - pu_\mu^{p-1}s}{s} = 0$$

and

$$\lim_{s \rightarrow \infty} \frac{(u_\mu + s)^p - u_\mu^p - pu_\mu^{p-1}s}{s^p} = 1.$$

Lemma 4.1. *For any $\epsilon > 0$, there exist a $C_\epsilon > 0$ such that*

$$(u_\mu + s)^p - u_\mu^p - pu_\mu^{p-1}s \leq \epsilon u_\mu^{p-1}s + C_\epsilon s^p$$

for all $s \geq 0$.

Lemma 4.2. *Suppose (H_1) with $\ell = -\frac{N+2}{2} - \epsilon$. There exists two constant $\alpha > 0$, $\rho > 0$ such that*

$$J_\mu(v) \geq \alpha > 0, \quad \text{for } v \in \mathcal{D}_0^1(\Omega), \quad \|v\| = \rho.$$

Proof. Lemma 4.1 implies that

$$\begin{aligned} J_\mu(v) &= \frac{1}{2} \int_\Omega |\nabla v|^2 dx - \frac{p}{2} \int_\Omega K(x) u_\mu^{p-1} (v^+)^2 dx \\ &\quad - \int_\Omega \int_0^{v^+} K(x) [(u_\mu + s)^p - u_\mu^p - pu_\mu^{p-1}s] ds dx \\ &\geq \frac{1}{2} \int_\Omega (|\nabla v|^2 - pK(x) u_\mu^{p-1} (v^+)^2) dx \\ &\quad - \int_\Omega K(x) \left(\frac{\epsilon}{2} u_\mu^{p-1} (v^+)^2 + C_\epsilon \frac{(v^+)^{p+1}}{p+1} \right) dx. \end{aligned}$$

Furthermore, from the definition of σ_μ in Lemma 3.2, we have

$$\int_\Omega |\nabla v|^2 dx \geq \sigma_\mu p \int_\Omega K(x) u_\mu^{p-1} (v^+)^2 dx$$

and, therefore, by $\sigma_\mu > 1$ we obtain by choosing ϵ small enough

$$\begin{aligned} J_\mu(v) &\geq \frac{1}{2\sigma_\mu} (\sigma_\mu - 1 - \epsilon) \int_\Omega |\nabla v|^2 dx - \frac{C_\epsilon}{p+1} \int_\Omega K(x) v^{p+1} dx \\ &\geq \frac{1}{4\sigma_\mu} (\sigma_\mu - 1) \int_\Omega |\nabla v|^2 dx - C \left[\int_\Omega |\nabla v|^2 dx \right]^{p+1} \\ &= \frac{1}{4\sigma_\mu} (\sigma_\mu - 1) \|v\|^2 - C \|v\|^{p+1} \end{aligned}$$

and the conclusion in Lemma 4.2 follows. \square

Lemma 4.3. *Suppose (H_1) with $\ell = -\frac{N+2}{2} - \epsilon$. Then there exist $0 < \psi_0 \in \mathcal{D}_0^1(\Omega)$ and $R_0 > 0$ such that*

$$J_\mu(R\psi_0) < 0$$

for $R \geq R_0$.

Proof. Let $h(x, s) = K(x)((u_\mu + s)^p - u_\mu^p - s^p)$, since $u_\mu(x)$ is bounded in Ω , it is easy to check that

$$\begin{aligned} \lim_{s \rightarrow 0} \frac{h(x, s)}{s} &\leq M \\ \lim_{s \rightarrow \infty} \frac{h(x, s)}{s^p} &= 0 \end{aligned}$$

uniformly in $x \in \Omega$, where $M > 0$ is some constant independent of x . Therefore, for any $\epsilon > 0$, there is a constant $C_\epsilon > 0$ such that

$$h(x, s) \leq \epsilon s^p + C_\epsilon s.$$

Now, choose a nonzero function $\psi_0 \in C_0^\infty(\Omega)$ such that $\psi_0 \geq 0$ and $K(x) \geq k_0 > 0$ on the support of ψ_0 . Then

$$J_\mu(R\psi_0) \leq \frac{1}{2}R^2\|\psi_0\|^2 - \frac{R^{p+1}}{p+1} \int_\Omega K(x)\psi_0^{p+1} dx + C_\epsilon R^2 \int_\Omega K\psi_0^2 dx + \epsilon R^{p+1} \int_\Omega K\psi_0^{p+1} dx.$$

It is then clear from the choice of ψ_0 , that for ϵ sufficiently small there is $R_0 > 0$ such that

$$J_\mu(R\psi_0) < 0 \text{ for all } R \geq R_0.$$

This completes the proof of Lemma 4.3, with R_0 and ψ_0 as above. \square

In order to use mountain pass Lemma [BN] to obtain the solution of $(4.1)_\mu$, we suppose moreover (H_2) .

Set

$$\Gamma = \{\gamma \in C([0, 1], \mathcal{D}_0^{1,2}(\Omega)), \gamma(0) = 0, \gamma(1) = R_0\psi_0\},$$

where ψ_0 is given by Lemma 4.2. We exploit the fact that the critical equation

$$-\Delta u = u^{\frac{N+2}{N-2}} \quad \text{in } \mathbb{R}^N$$

has the positive radial solution

$$u_\epsilon(x) = k \left[\frac{\epsilon}{\epsilon^2 + |x - x_0|^2} \right]^{\frac{N-2}{2}}$$

with $k = (N(N-2))^{\frac{N-2}{4}}$ for any $\epsilon > 0$, $x \in \mathbb{R}^N$. Furthermore,

$$\int_{\mathbb{R}^N} |\nabla u_\epsilon|^2 dx = \int_{\mathbb{R}^N} u_\epsilon^{p+1} dx = S^{N/2},$$

and for some positive constant c depending only on N $cu_\epsilon(x)$ attains the infimum for the variational problem

$$S = \inf \left\{ \|u\|^2 \mid \int_{\mathbb{R}^N} u^{p+1} dx = 1 \quad u \in \mathcal{D}_0^{1,2}(\mathbb{R}^N) \right\}.$$

Let $R > 0$ be small enough that $B_{2R}(x_0) \in V$. Let ψ be a piecewise smooth function with support in B_{2R} such that $\psi(x) \equiv 1$ in $B_R(x_0)$, $0 \leq \psi(x) \leq 1$ in $B_{2R}(x_0)$ and $|\nabla \psi(x)| \leq \frac{1}{R}$. Define

$$w_\epsilon(x) = \psi(x)u_\epsilon(x)$$

and

$$v_\epsilon(x) = w_\epsilon(x) \left[\int_{\Omega} K(x)w_\epsilon^{p+1} dx \right]^{\frac{-1}{p+1}}.$$

The proof of the following Lemma follows the same lines as in [BK].

Lemma 4.4. *If assumptions (H_1) - (H_2) holds and $p = \frac{N+2}{N-2}$, then there exist some positive constant $\epsilon > 0$ and $t_0 > 0$ such that*

$$J_\mu(t_0 v_\epsilon) < 0$$

and

$$0 < \sup_{t \geq 0} J_\mu(t v_\epsilon) < \frac{1}{N} S^{\frac{N}{2}} (\|K\|_{L^\infty})^{\frac{2-N}{2}}.$$

Proof. Since $\frac{\partial u_\epsilon}{\partial \gamma} \leq 0$, we have

$$\int_{B_R} |\nabla w_\epsilon|^2 dx = \int_{B_R} |\nabla u_\epsilon|^2 dx \leq \int_{B_R} u_\epsilon^{p+1} dx$$

and by the assumption (H₂) we also have

$$K(x_0) \int_{B_R} u_\epsilon^{p+1} dx \leq \int_{B_R} K(x) u_\epsilon^{p+1} dx + 0(\epsilon^2).$$

Simple calculations also show that

$$\begin{aligned} \int_{\mathbb{R}^N \setminus B_R} u_\epsilon^{p+1} dx &= 0(\epsilon^N) \\ A_\epsilon &\equiv \int_{\mathbb{R}^N \setminus B_R} |\nabla w_\epsilon|^2 dx = 0(\epsilon^{N-2}) \end{aligned}$$

as $\epsilon \rightarrow 0$ and

$$S = \left[\int_{\mathbb{R}^N} u_\epsilon^{p+1} dx \right]^{\frac{2}{N}}.$$

Therefore, we have

$$\begin{aligned} \int_{\mathbb{R}^N} |\nabla w_\epsilon|^2 dx &= \int_{B_R} |\nabla w_\epsilon|^2 dx + A_\epsilon \\ &\leq \int_{B_R} u_\epsilon^{p+1} dx + A_\epsilon \\ &\leq S \left[\int_{B_R} u_\epsilon^{p+1} dx \right]^{\frac{2}{p+1}} + A_\epsilon \\ &\leq S \|K\|_\infty^{-\frac{2}{p+1}} \left[\int_{B_R} K(x) w_\epsilon^{p+1} dx \right]^{\frac{2}{p+1}} + 0(\epsilon^2) + 0(\epsilon^{N-2}). \end{aligned}$$

Set $V_\epsilon \equiv \int_{\mathbb{R}^N} |\nabla v_\epsilon|^2 dx$, since for small $\epsilon > 0$, say $\epsilon \leq \epsilon_0$, it is easy to see that

$$\int_{B_R} K(x) w_\epsilon^{p+1} dx \geq C_{\epsilon_0}$$

for some positive constant C_{ϵ_0} , the definition of V_ϵ and the last two inequalities imply that

$$V_\epsilon \leq S(\|K\|_\infty)^{\frac{2}{p+1}} + 0(\epsilon^2) + 0(\epsilon^{N-2}).$$

We consider now $J_\mu(v)$

$$\begin{aligned} J_\mu(v) &= \frac{1}{2} \int_{\Omega} |\nabla v|^2 dx - \frac{1}{p+1} \int_{\Omega} K(x) [(u_\mu + v^+)^{p+1} - u_\mu^{p+1}] dx + \int_{\Omega} K(x) u_\mu^p v^+ dx \\ &= \frac{1}{2} \int_{\Omega} |\nabla v|^2 dx - \int_{\Omega} K(x) \int_0^{v^+} ((u_\mu + s)^p - u_\mu^p) ds dx \\ &= \frac{1}{2} \int_{\Omega} |\nabla v|^2 dx - \frac{1}{p+1} \int_{\Omega} K(x) (v^+)^{p+1} dx \\ &\quad - \int_{\Omega} K(x) \int_0^{v^+} [(u_\mu + s)^p - u_\mu^p - s^p] ds dx. \end{aligned}$$

Set $F(x, v) = K(x) \int_0^{v^+} ((u_\mu + s)^p - u_\mu^p - s^p) ds$, then

$$J_\mu(tv_\epsilon) = \frac{1}{2}t^2V_\epsilon - \frac{1}{p+1}t^{p+1} - \int_\Omega F(x, tv_\epsilon)dx .$$

Clearly, $\lim_{t \rightarrow \infty} J_\mu(tv_\epsilon) = -\infty$ for all $\epsilon > 0$, hence $\sup_{t \geq 0} J_\mu(tv_\epsilon)$ is attained by some $t_\epsilon \geq 0$, we may assume $t_\epsilon > 0$ for $\epsilon > 0$, otherwise there would be nothing to prove.

It follows from $\frac{d}{dt}J_\mu(tv_\epsilon)|_{t=t_\epsilon} = 0$ and the monotonicity of F in v that

$$t_\epsilon \leq V_\epsilon^{\frac{1}{p-1}} \leq C_0 \text{ for all } \epsilon > 0 ,$$

where C_0 is some positive constant independent of ϵ . By the monotonicity property of $\frac{1}{2}t^2V_\epsilon - \frac{1}{p+1}t^{p+1}$ on the interval $(0, V_\epsilon^{\frac{1}{p-1}}]$ we then have

$$\sup_{t \geq 0} J_\mu(tv_\epsilon) = J_\mu(t_\epsilon v_\epsilon) \leq \frac{1}{N}V_\epsilon^{\frac{N}{2}} - \int_{B_{2R}} F(x, t_\epsilon v_\epsilon)dx .$$

The estimate on V_ϵ and the above inequality imply that

$$\sup_{t \geq 0} J_\mu(tv_\epsilon) \leq \frac{1}{N}S^{\frac{N}{2}}(\|k\|_\infty)^{\frac{2-N}{2}} - \int_{B_{2R}} F(x, tv_\epsilon)dx + 0(\epsilon^L) ,$$

where $L = \min(N-2, 2)$. The conclusion will follow if we can show that

$$\lim_{\epsilon \rightarrow 0} \epsilon^{-L} \int_{B_{2R}} F(x, t_\epsilon v_\epsilon)dx = +\infty .$$

First we claim that

$$\lim_{\epsilon \rightarrow 0} t_\epsilon > 0 .$$

Indeed, by $\frac{d}{dt}J_\mu(tv_\epsilon)|_{t=t_\epsilon} = 0$ we have

$$V_\epsilon - t_\epsilon^{p-1} - t_\epsilon^{-1} \int_\Omega K(x)[(u_\mu + t_\epsilon v_\epsilon)^p - u_\mu^p - t_\epsilon^p v_\epsilon^p]v_\epsilon dx = 0 .$$

We show that

$$\lim_{\epsilon \rightarrow 0} t_\epsilon^{-1} \int_\Omega K(x)[(u_\mu + t_\epsilon v_\epsilon)^p - u_\mu^p - t_\epsilon^p v_\epsilon^p]v_\epsilon dx = 0 .$$

This will follow by the same procedure as in [BK] (p465–466) by observing first that for all $\delta > 0$, $\exists C_\delta > 0$ such that

$$|f(x, u)| \equiv |(u_\mu + u)^p - u_\mu^p - u^p| \leq \delta u^p + C_\delta u ,$$

for all $u > 0$. This follows easily from the boundedness of u_μ . Indeed for $u \geq \frac{1}{\delta}$, we have

$$|f(x, u)| = u^p p \int_0^{\frac{u_\mu}{u}} ((s+1)^{p-1} - s^{p-1}) ds \leq C_\delta u^p$$

for some constant C . For $u \leq \frac{1}{\delta}$ we have

$$\begin{aligned} |f(x, u)| &\leq \left| \frac{(u_\mu + u)^p - u_\mu^p}{u} \right| u + u^p \\ &\leq p(u_\mu + u)^{p-1} u + \left(\frac{1}{\delta} \right)^{p-1} \\ &\leq C_\delta u . \end{aligned}$$

It then follows that a positive constant C , independent of ϵ , exists such that

$$\sup_{t \geq 0} J_\mu(tv_\epsilon) \leq \frac{1}{N} S^{\frac{N}{2}} (\|K\|_\infty)^{\frac{2-N}{2}} - \int_{B_{2R}} F(x, Cv_\epsilon) dx + o(\epsilon^L)$$

for sufficiently small $\epsilon > 0$. A change of variables yields

$$\lim_{\epsilon \rightarrow 0^+} \epsilon^{-L} \int_{B_{2R}} F(x, Cv_\epsilon) dx = +\infty$$

as in [BN]. □

Lemma 4.5. *Assume H_2) and H_1). Suppose moreover $0 \not\equiv K(x) \geq 0$ and $K(x) \in L^1(\Omega)$. Then problem (4.1) $_\mu$ has at least two solution for each $\mu \in (0, \mu^*)$ if $p = \frac{N+2}{N-2}$.*

Proof. The conditions for the mountain pass Lemma [BN] are satisfied by Lemma 4.2, 4.3. Hence there is a sequence $\{v_n\} \subset \mathcal{D}_0^1(\Omega)$ such that $J_\mu(v_n) \rightarrow c$ and $J'_\mu(v_n) \rightarrow 0$ in $\mathcal{D}_0^1(\Omega)$ as $n \rightarrow \infty$, where

$$c = \inf_{\nu \in \Gamma} \sup_{u \in \nu} J_\mu(u) .$$

Thus

$$J_\mu(v_n) = \frac{1}{2} \int_\Omega |\nabla v_n|^2 dx - \int_\Omega \left[\frac{1}{p+1} K(x) (u_\mu + v_n^+)^{p+1} - \frac{1}{p+1} u_\mu^{p+1} - u_\mu^p v_n^+ \right] dx = c + o(1), \quad (4.2)$$

and

$$J'_\mu(v_n)\psi = \int_\Omega \nabla v_n \nabla \psi dx - \int_\Omega K((u_\mu + v_n^+)^p - u_\mu^p) \psi dx = o(1) \|\psi\| \quad (4.3)$$

as $n \rightarrow \infty$ and $\psi \in \mathcal{D}_0^1(\Omega)$. Choose $\frac{1}{p+1} < \theta < \frac{1}{2}$ and $\psi = v_n$. It follows from (4.2), (4.3) that

$$\begin{aligned}
c + o(1) &\geq \frac{1}{2} \int_{\Omega} |\nabla v_n|^2 dx - \frac{1}{p+1} \int_{\Omega} K(u_{\mu} + v_n^+)^{p+1} dx \\
&= \left(\frac{1}{2} - \theta\right) \|v_n\|^2 + \theta(\|v_n\|^2 - \int_{\Omega} K[(u_{\mu} + v_n^+)^p - u_{\mu}^p]v_n dx) \\
&\quad + \left(\theta - \frac{1}{p+1}\right) \int_{\Omega} K(x)(u_{\mu} + v_n^+)^p v_n^+ dx - \theta \int_{\Omega} K u_{\mu}^p v_n^+ dx \\
&\quad - \frac{1}{p+1} \int_{\Omega} K u_{\mu} (u_{\mu} + v_n^+)^p dx \\
&= \left(\frac{1}{2} - \theta\right) \|v_n\|^2 + o(1)\|v_n\| \\
&\quad + \left(\theta - \frac{1}{p+1}\right) \int_{\Omega} K(x)(v_n^+ - \tau u_{\mu})(u_{\mu} + v_n^+)^p dx \\
&\quad - \theta \int_{\Omega} K(x)u_{\mu}^p v_n^+ dx,
\end{aligned}$$

where $\tau = (p+1)^{-1}(\theta - (p+1)^{-1})^{-1}$. Notice that we have used the obvious equality

$$\int_{\Omega} K(x)((u_n + v_n^+)^p - u_n^p)v_n dx = \int_{\Omega} K[(u_{\mu} + v_n^+)^p - u_{\mu}^p]v_n^+ dx.$$

Using Hölder's and Sobolev's inequality we have

$$\begin{aligned}
\int_{\Omega} K u_{\mu}^p v_n^+ dx &\leq \|u_{\mu}\|_{\infty}^p \int_{\Omega} K(x)v_n^+ dx \\
&\leq C \left(\int_{\Omega} K(x)^2 dx \right)^{\frac{1}{2}} \left(\int v_n^{+2} dx \right)^{\frac{1}{2}} \\
&\leq C_1 \|v_n^+\|
\end{aligned}$$

for some constant $C > 0$.

Because $g(x) = (s - \tau u_{\mu})(u_n + s)^p$ gets its minimum at

$$s = \frac{p\tau - 1}{1 + p}$$

we have

$$\begin{aligned}
c + o(1) &\geq \left(\frac{1}{2} - \theta\right) \|v_n\|^2 + o(1)\|v_n\| \\
&\quad - \frac{2(p(1 + \tau))^p(\theta(p+1) - 1)}{(p+1)^{p+2}} \int_{\Omega} K(x)u_{\mu}^{p+1} dx \\
&\quad - \theta C_1 \|v_n\|
\end{aligned}$$

since $\|u_\mu\|_{L^\infty}$ is bounded, $K(x) \in L^1(\Omega)$. From the above inequality we can deduce $\{v_n\}$ is bounded in $\mathcal{D}_0^1(\Omega)$. Standard embedding theorem then show that $\{v_n\}$ has a subsequence, still denoted by $\{v_n\}$ for which

$$\begin{aligned} v_n &\rightharpoonup v \text{ weakly in } \mathcal{D}_0^1(\Omega) \\ v_n &\rightarrow v \text{ a.e. in } \Omega \\ v_n &\rightharpoonup v \text{ weakly in } L^{p+1}(\Omega). \end{aligned}$$

It follows from (4.2) and (4.3) that v is a weak solution of

$$-\Delta v = K(x)[(u_\mu + v^+)^p - u_\mu^p] \quad v \in \mathcal{D}_0^1(\Omega).$$

Furthermore, (4.3) with $\psi = v^-$ implies that $\int_\Omega |\nabla v^-|^2 dx = 0$ and therefore $\int_\Omega |v^-|^{p+1} dx = 0$, by Sobolev embedding. This shows that $v \geq 0$ a.e. in Ω , we show next that $v \not\equiv 0$.

Consider the sequence $\{w_n\}$, $w_n = v_n - v$, for a subsequence of $\{w_n\}$, denoted the same way, we define

$$\ell = \lim_{n \rightarrow \infty} \|w_n\|^2.$$

If $\ell = 0$, the continuity of J_μ on $\mathcal{D}_0^1(\Omega)$ implies that

$$0 < \alpha \leq c = \lim_{n \rightarrow \infty} J_\mu(v_n) = J_\mu(v) \quad \text{and hence } v \not\equiv 0.$$

If $\ell > 0$, we proceed as follows. Using (4.3) with $\psi = v_n$, the boundedness of $\|v_n\|$, the weak convergence of v_n to v in $L^{p+1}(\Omega)$ and the fact that $u_\mu \in L^\infty(\Omega)$, $K(x) \in L^{\frac{p+1}{p}}$ we obtain

$$\begin{aligned} \int_\Omega K(x)(u_\mu + v_n)^p u_\mu dx &\rightarrow \int_\Omega K(x)(u_\mu + v)^p u_\mu dx \\ \int_\Omega K(x)u_\mu^p v_n dx &\rightarrow \int_\Omega K(x)u_\mu^p v dx \quad . \end{aligned}$$

We have

$$\int_\Omega |\nabla v_n|^2 dx - \int_\Omega K(x)(u_\mu + v_n^+)^{p+1} dx + \int_\Omega K(x)(u_\mu + v)^p u_\mu dx + \int_\Omega K(x)u_\mu^p v dx = o(1). \quad (4.4)$$

Using a lemma of Brezis and Lieb [BL] and (4.4) we obtain

$$\begin{aligned} \int_\Omega |\nabla w_n|^2 dx + \int_\Omega |\nabla v|^2 dx - \int_\Omega K(x)(v_n^+ - v)^{p+1} dx \\ = \int_\Omega K(x)((u_\mu + v)^p - u_\mu^p)v dx + o(1). \end{aligned}$$

Since v is a solution of problem $(4.1)_\mu$, we have

$$\int_{\Omega} |\nabla w_n|^2 dx = \int_{\Omega} K(x)(v^+ - v)^{p+1} dx + o(1). \quad (4.5)$$

Using (4.3) and Brezis-Lieb Lemma [BL] we also have

$$\begin{aligned} o(1) + c &= \frac{1}{2} \int_{\Omega} |\nabla w_n|^2 dx + \frac{1}{2} \int_{\Omega} |\nabla v|^2 dx - \frac{1}{p+1} \int_{\Omega} K(x) w_n^{p+1} dx \\ &\quad - \frac{1}{p+1} \int_{\Omega} K(x) (u_\mu + v)^{p+1} dx + \frac{1}{p+1} \int_{\Omega} K(x) u_\mu^{p+1} dx + \int_{\Omega} K(x) u_\mu^p v dx \\ &= \frac{1}{2} \int_{\Omega} |\nabla w_n|^2 dx - \frac{1}{p+1} \int_{\Omega} K(x) (v_n^+ - v)^{p+1} dx + J_\mu(v). \end{aligned}$$

which gives us

$$o(1) + c = \frac{1}{2} \int_{\Omega} |\nabla w_n|^2 dx - \frac{1}{p+1} \int_{\Omega} K(x) (v_n^+ - v)^{p+1} dx + J_\mu(v). \quad (4.6)$$

It follows from (4.5) and (4.6) that

$$c = \frac{1}{N} \ell + J_\mu(v).$$

We also have by Sobolev's inequality and (4.5) that

$$\begin{aligned} \int_{\Omega} |\nabla w_n|^2 dx &\geq S \left(\int_{\Omega} |w_n|^{p+1} dx \right)^{\frac{2}{p+1}} \\ &\geq S \left(\int_{\Omega} |v_n^+ - v|^{p+1} dx \right)^{\frac{2}{p+1}} \\ &\geq \left(\frac{1}{\sup_{\Omega} K(x)} \right)^{\frac{2}{p+1}} S \left(\int_{\Omega} K(x) (|v_n^+ - v|^{p+1}) dx \right)^{\frac{2}{p+1}} \\ &= \left(\frac{1}{\sup_{\Omega} K} \right)^{\frac{2}{p+1}} S (\|w_n\|^2 + o(1))^{\frac{1}{p+1}} \end{aligned}$$

which gives in the limit, as $n \rightarrow \infty$, the inequality

$$\ell \geq [\sup_{\Omega} K(x)]^{-\frac{2}{p+1}} S \ell^{\frac{2}{p+1}} \quad (4.8)$$

since $\ell > 0$, (4.7) and (4.8) give

$$c \geq \frac{1}{N} (\sup_{\Omega} K(x))^{\frac{2-N}{2}} S^{\frac{N}{2}} + J_\mu(v) \quad (4.9)$$

which implies from Lemma 4.4 that $J_\mu(v) < 0$, thus $v \not\equiv 0$. \square

From the above lemmas we conclude the theorem 1.2.

References

- [B] V. Benci and G. Cerami, Positive solutions of some nonlinear elliptic problems in exterior domains, *Arch. Rat'l. Mech. Anal.* **99** (1987) pp. 283-300.
- [BK] H. Brezis and T. Kato, Remarks on the Schrodinger operator with singular complex potentials, *J. Math. Pures. Appl.* **58** (1979), 137-151.
- [BLY] R. Benguria, S. Lorca and C. Yarur, Nonexistence results for solutions of semilinear elliptic equations, *Duke Math. J.* **74** (1994), 615-634.
- [BL] H. Brezis and E. Lieb, A relation between pointwise convergence of function and convergence of functional, *Proc. Amer. Math. Soc.* **88** (1993) pp. 486-490.
- [BN] H. Brezis and L. Nirenberg, Positive solutions of nonlinear elliptic equations involving critical Sobolev exponents, *Comm. Pure. Appl. Math.* **36** (1983) pp. 437-477
- [CZ] D.-M. Cao and H.-S. Zhou, Multiple positive solution of nonhomogeneous semilinear elliptic equations in \mathbb{R}^N , *Proc. Royal Society Edinburgh* **126A** (1996) 443-463.
- [CL1] W.-X. Chen and C.-M Li, A necessary and sufficient condition for the Nirenberg problem, *Comm. Pure Appl. Math.* **48** (1995), 657-667.
- [CL2] W.-X. Chen and C.-M Li, A priori estimates for prescribing scalar curvature equations, *Ann. of Math.* **145** (1997), 547-564.
- [DL1] Y.-B. Deng and Y. Li, Existence and Bifurcation of the Positive Solutions For a Semilinear Equation With Critical Exponent, *J. Diff. Equns.* **130(1)** (1996) 179-200.
- [DL2] Y.B. Deng and Y. Li, Existence and of Multiple Positive Solutions For a Semilinear Elliptic Equation *Advances Diff. Equns.* **2(3)** (1997), 361-382.
- [DLZ] Y.-B. Deng, Y. Li and X.-J. Zhao, Multiple solutions for an inhomogeneous semilinear elliptic equation in \mathbb{R}^n , preprint.
- [DN1] W.-Y. Ding and W.-M. Ni, On the elliptic equation $\Delta u + Ku^{(n+2)/(n-2)} = 0$ and related topics, *Duck Math. J.* **52**, 485-506(1985),

- [DN2] W.-Y. Ding and W.-M. Ni, On the existence of positive entire solution of a semilinear elliptic equation, *Arch. Rat'l Mech. Anal.* **91** (1986) pp. 283-308.
- [E] H. Egnell, Asymptotic results for finite energy solutions of semilinear elliptic equations, *J. Diff. Equ.* **98** (1992), No.1, 34-58.
- [Es] J. Escobar, Positive solutions for some semilinear elliptic equations with critical Sobolev exponents, *Comm. Pure and Appl. Math.*, , (1987), 623-657.
- [G] C.-F. Gui, Entire Positive Solutions of Equation $\Delta u + F(x, u) = 0$ J.D.E., 1993
- [GE] J. Graham-Eagle, Monotone method for semilinear elliptic equations in unbounded domains, *J. Math. Anal. Appl.* **137** (1989), 122-31.
- [JPY] R.A. Johnson, X.-B. Pan and Y.-F. Yi, Positive Solutions of Super-Critical Elliptic Equations and Asymptotics, *Comm. Partial Diff. Eqns*, 18(5-6), 977-1019(1993).
- [KN] C. E. Kening and W.-M. Ni, An exterior Dirichlet problem with applications to some nonlinear equations arising in geometry, *Amer. J. Math.* **106**, 689-702 (1984)
- [KL] M.-K. Kwong and Y. Li, Uniqueness of radial solutions of semilinear elliptic equations, *Trans. Amer. Math. Soc.*, 333(1992), 339-363.
- [L] Y. Li, Asymptotic Behavior of Positive Solutions of Equations $\Delta u + K(x)u^p = 0$ in R^n , *J.D.E.* 95(1992), 304-330.
- [LN1] Y. Li and W.-M. Ni, On Conformal Scalar Curvature Equations in R^n , *Duck Math. J.* 57 895-924(1988).
- [LN2] Y. Li and W.-M. Ni, On the asymptotic behavior and radial symmetry of positive solutions of semilinear elliptic equations in R^N , *Arch. Rational Mech. Anal.* **118** (1992) 195-222.
- [LY1] Y.-Y. Li, Prescribing scalar curvature on S^n and related problems. I., *J. Differential Equations* **120** (1995), 319-410.
- [LY2] Y.-Y. Li, Prescribing scalar curvature on S^n and related problems. II. Existence and compactness, *Comm. Pure Appl. Math.* **49**(1996), 541-597.

- [Lin] F.-H. Lin, On the elliptic equation $D_i[a_{ij}(x)D_jU] - k(x)U + K(x)U^p = 0$, *Proc. Amer. Math. Soc.* **95** (1985), 219-226.
- [Lio] P. L. Lions, The concentration-compactness principle in the calculus of variations, *Revista Matemática Iberoamericana* **1(2)** (1985), 45-121.
- [N1] W.-M. Ni, On the elliptic equation $\Delta u + K(x)u^{(n+2)/(n-2)} = 0$, it's generalizations and applications in geometry, *Indiana Univ. Math J.* 31,493-529(1982).
- [N2] W.-M. Ni, Some aspects of semilinear elliptic equations in R^n , in "Nonlinear Diffusion Equations and Their Equilibrium States" (W.-M. Ni, L.A. Peletier, and J. Serrin, Eds.), Springer-Verlag, New York, 1988.
- [NY] W.-M. Ni and S. Yotsutani, Semilinear elliptic equations of Matukuma type and related topics, *Japan J. Appl. Math.* 5 (1988), 1-32.
- [WW] X.-F. Wang and J.-C. Wei, On the equation $\Delta u + K(x)u^{(n+2)/(n-2)\epsilon^2} = 0$ in R^n , *Rend. Circ. Mat. Palermo* **44** (1995), 365-400.
- [Y] J.-F. Yang, Positive solution of semilinear elliptic problems in exterior domains, *J. Diff. Equa.* **106** (1993) pp.40-69.
- [YY] E. Yanogida and S. Yosutani, Classification of positive radial solutions to $\Delta u + K(|x|)u^p = 0$ in \mathbf{R}^n , *Arch. Rational Mech. Anal.*, 124(1993),239-259.
- [Z] Z. Zhao, On the existence of positive solution of nonlinear elliptic equations—a probabilistic potential theory approach, *Duke Math. J.* **69 (2)** (1993), 247-258.
- [ZC] X.-P. Zhou and D.-M. Cao, The concentration-compactness principle nonlinear elliptic equations, *Acta Math.Sci.* **9** (1989) 307-28.