

$T_p(M) = \{v : G(p) \rightarrow \mathbf{R} \mid v \text{ is linear and satisfies the Leibniz rule } \}$

$G(p) = \{[g] \mid g^{\text{smooth}} : U \rightarrow \mathbf{R}, \text{ for some } U^{\text{open}} \text{ such that } p \in U \subset M\}$  is an algebra over  $\mathbf{R}$ .

$v \in T_p(M)$  is called a *derivation*

The directional derivative of  $[g]$  in direction  $[\alpha] =$

$$D_\alpha g = \frac{d(g \circ \alpha)}{dt} \Big|_{t=0} = g'(\alpha(0))\alpha'(0) \in \mathbf{R}$$

Properties:

$D_\alpha$  is linear and satisfies the Leibniz rule. i.e,

$$1a.) D_\alpha(g + h) = D_\alpha g + D_\alpha h \qquad 1b.) D_\alpha(cg) = cD_\alpha g$$

$$2.) D_\alpha(g \cdot h) = D_\alpha g \cdot h(p) + g(p) \cdot D_\alpha h$$

Thus  $D_\alpha \in T_p(M)$

Thm: Let  $M$  be an  $m$ -manifold, then  $T_p(M)$  is an  $m$ -dimensional real vector space.  $[(c_1v + c_2w)(f) = c_1v(f) + c_2w(f)]$ .

Take a chart  $(U, \phi)$  at  $p$  where  $\phi(p) = \mathbf{0}$ ,

The *standard basis* for  $T_p(M)$  w.r.t.  $(U, \phi) = \{v_1, \dots, v_m\}$ ,

where  $v_i = D_{\alpha_i}$  and

$$\alpha_i : (-\epsilon, \epsilon) \rightarrow M, \alpha_i(t) = \phi^{-1}(0, \dots, t, \dots, 0) \text{ for some } \epsilon > 0.$$

Prop:  $\{v_1, \dots, v_m\}$  are linearly independent.

Proof: Evaluate  $v_i$  at a “projection map”.

Thm:  $\{v_1, \dots, v_m\}$  span  $T_p(M)$ .

If  $v \in T_p(M)$ , then  $v = \sum_{i=1}^m a_i v_i$  where  $a_i = v([\pi_i \circ \phi])$

Prop: If  $v$  is a derivation, and  $f$  is constant, then  $v(f) = 0$ .

Suppose  $f^{\text{smooth}} : M \rightarrow N, f(p) = q$ .

The *tangent (or differential) map*,

$$df_p : T_p M \rightarrow T_q N$$

$df_p(v)$  is the derivation which takes  $[g] \in G(q)$  to the real number  $v([g \circ f])$

I.e.,  $df_p$  takes the derivation  $(v : G(p) \rightarrow \mathbf{R}) \in T_p M$

to the derivation in  $T_q N$  which takes

the germ  $[g] \in G(q)$  to the real number  $v([g \circ f])$ .

Suppose  $f : M \rightarrow N$  is smooth. Then  $f^* : G(q) \rightarrow G(p)$  is a homomorphism.

Let  $V$  be a real vector space.

Then the dual space,  $V^* = \{f : V \rightarrow \mathbf{R} \mid f \text{ is linear } \}$

Proposition: With respect to this choice of basis, the matrix of  $d_p f : T_p M \rightarrow T_q N$  is  $(\frac{\partial F_i}{\partial x_j})_{\varphi(p)}$ , where  $x_j$  are coordinates in  $\mathbf{R}^m$  and  $F_i = (F_1, \dots, F_n)$  in coordinates for  $\mathbf{R}^n$ .

Here are some more properties of  $d_p f$  and  $T_p M$ : Let  $f: M \rightarrow N$  be smooth.

- 1.) If  $f: M \rightarrow N$  is a diffeomorphism, the  $d_p f$  is an isomorphism, for all  $p \in M$ .
- 2.) If  $d_p f = 0$  for all  $p \in M$  iff  $f$  is a constant map.
- 3.) If  $id: M \rightarrow M$ ,  $id(x) = x$ , then  $d_p(id) = I_m$
- 4.)  $d_r(f \circ h) = d_p(f) \circ d_r(h)$  where  $p = h(r)$ .
- 5.) If  $N \cong M/G$ , where  $G$  is a discrete Lie group acting properly discontinuously on  $M$ , and  $f: M \rightarrow M/G$  is the orbit map, the  $d_p f$  is an isomorphism for all  $p$ .
- 6.)  $T_{(p,q)}(M \times N) \cong T_p(M) \times T_q(N)$ .

Rank Theorem: Suppose  $A_0 \subset \mathbf{R}^n$ ,  $B_0 \subset \mathbf{R}^m$ ,  $F: A_0 \rightarrow B_0 \in C^1$   $a \in A_0, b \in B_0$ . Suppose  $\text{rank } F = k$ .

Then there exists  $A^{open} \subset A_0$  such that  $a \in A$  and  $B^{open} \subset B_0$  such that  $b \in B$  and  $G, H, C^r$  diffeomorphisms

such that  $G: A \rightarrow U^{open} \subset \mathbf{R}^n$ ,  $H: B \rightarrow V^{open} \subset \mathbf{R}^m$  and

$$H \circ F \circ G^{-1}(x_1, \dots, x_n) = (x_1, \dots, x_k, 0, \dots, 0)$$

---

Proposition:  $\text{rank } DF(a) = k$  implies there exists  $V$  open such that  $a \in V$  and  $DF(x) \geq k$  for all  $x \in V$

Proof:

Rank of  $A = \dim(\text{span}\{\mathbf{r}_1, \dots, \mathbf{r}_m\}) = \dim(\text{span}\{\mathbf{c}_1, \dots, \mathbf{c}_m\})$   
 = maximum order of any nonvanishing minor determinant.

Use determinant is a continuous function.

---

Thm 6.4 (Inverse Function Theorem): Suppose  $F: W^{open} \subset \mathbf{R}^n \rightarrow \mathbf{R}^n \in C^r$ . Suppose for  $a \in W$ ,  $\det(DF_a) \neq 0$ . Then there exists  $U$  such that  $a \in U^{open}$ ,  $V = F(U)$  is open, and  $F: U \rightarrow V$  is a  $C^r$ -diffeomorphism. Moreover, for  $x \in U$  and  $y = F(x)$ ,  $DF_y^{-1} = (DF_x)^{-1}$