**Problem 1.** Suppose A is a  $5 \times 6$  matrix. If rank of A = 4, then nullity of A = 4

G. 2

## Problem 2.

Let 
$$A = \begin{bmatrix} 1 & -1 \\ 1 & 3 \end{bmatrix}$$
. Is  $A =$  diagonalizable?

• B. no

**Problem 3.** Suppose A is a 3  $\times$  4 matrix. Then *nul A* is a subspace of  $R^k$  where k =

• I. 4

**Problem 4.** Suppose  $A = PDP^{-1}$  where D is a diagonal matrix. Suppose also the  $d_{ii}$  are the diagonal entries of D. If  $P = [\vec{p_1} \ \vec{p_2} \ \vec{p_3}]$  and  $d_{11} = d_{33}$ , then  $\vec{p_1} + \vec{p_3}$  is an eigenvector of A

• A. True

## Problem 5.

Which of the following is an eigenvalue of  $\begin{bmatrix} 4 & 4 \\ 1 & 4 \end{bmatrix}$ .

• G. 2

**Problem 6.** If  $\vec{x_1}$  and  $\vec{x_2}$  are solutions to  $A\vec{x} = \vec{b}$ , then  $-5\vec{x_1} + 8\vec{x_2}$  is also a solution to  $A\vec{x} = \vec{b}$ .

• B. False

## Problem 7.

Calculate the determinant of  $\begin{bmatrix} -1.125 & -1 \\ 8 & 8 \end{bmatrix}$ .

• D. -1

Problem 8.

Suppose the orthogonal projection of  $\begin{bmatrix} -4 \\ 7 \\ 2 \end{bmatrix}$  onto  $\begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix}$  is  $(z_1, z_2, z_3)$ . Then  $z_1 =$ 

• B. -3

**Problem 9.** Suppose A is a square matrix and  $A\vec{x} = \vec{0}$  has an infinite number of solutions, then given a vector  $\vec{b}$  of the appropriate dimension,  $A\vec{x} = \vec{b}$  has

• E. either no solution or an infinite number of solutions

Problem 10.

Let 
$$A = \begin{bmatrix} 8 & -24 & 32 \\ 0 & 2 & 8 \\ 0 & 0 & 8 \end{bmatrix}$$
. Is  $A = \text{diagonalizable}$ ?

• A. yes

## Problem 11.

$$\operatorname{Let} A = \begin{bmatrix} 5.31034482758621 & 2.12413793103448 & -5.7448275862069 \\ 4.22413793103448 & 1.68965517241379 & -5.7448275862069 \\ 0 & 0 & -2.46206896551724 \end{bmatrix}$$

and let 
$$P = \begin{bmatrix} -2 & -4 & 7 \\ 5 & -7 & 7 \\ 0 & -8 & 3 \end{bmatrix}$$
.

Suppose  $A = PDP^{-1}$ . Then if  $d_{ii}$  are the diagonal entries of D,  $d_{11} =$ 

• E. 0

**Problem 12.** Suppose  $\begin{bmatrix} u_1 \\ u_2 \\ u_3 \end{bmatrix}$  is a unit vector in the direction of  $\begin{bmatrix} 5 \\ 2 \\ 3.17214438511238 \end{bmatrix}$ . Then  $u_1 =$ 

• I. 0.8

**Problem 13.** Suppose  $A \begin{bmatrix} -2 \\ -2 \\ -1 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$ . Then an eigenvalue of A is

• E. 0

**Problem 14.** The vector  $\vec{b}$  is in *ColA* if and only if  $A\vec{v} = \vec{b}$  has a solution

• A. True

Problem 15.

Let 
$$A = \begin{bmatrix} 15 & -6 \\ 5 & -2 \end{bmatrix}$$
.

Which of the following could be a basis for null(A)?

• B. 
$$\left\{ \begin{bmatrix} 2 \\ 5 \end{bmatrix} \right\}$$