

4. Cancès, E., Mennucci, B.: New applications of integral equations methods for solvation continuum models: ionic solutions and liquid crystals. *J. Math. Chem.* **23**, 309–326 (1998)
5. Cancès, E., Le Bris, C., Mennucci, B., Tomasi, J.: Integral equation methods for molecular scale calculations in the liquid phase. *Math. Models Methods Appl. Sci.* **9**, 35–44 (1999)
6. Cortis, C., Friesner, R.: An automatic three-dimensional finite element mesh generation system for the Poisson–Boltzmann equation. *J. Comput. Chem.* **18**(13), 1570–1590 (1997)
7. Gao, J.L.: Hybrid quantum and molecular mechanical simulations: an alternative avenue to solvent effects in organic chemistry. *Acc. Chem. Res.* **29**(6), 298–305 (1996)
8. Holst, M., Baker, N., Wang, F.: Adaptive multilevel finite element solution of the Poisson–Boltzmann equation I. algorithms and examples. *J. Comput. Chem.* **21**, 1319–1342 (2000)
9. Klamt, A.: The COSMO and COSMORS solvation models. *Wiley Interdiscip. Rev. Comput. Mol. Sci.* **1**(5), 699–709 (2011)
10. Lin, H., Truhlar, D.G.: QM/MM: What have we learned, where are we, and where do we go from here? *Theor. Chem. Acc.* **117**, 185–199 (2007)
11. Mennucci, B.: Continuum solvation models: what else can we learn from them? *J. Phys. Chem. Lett.* **1**(10), 1666–1674 (2010)
12. Senn, H.M., Thiel, W.: QM/MM methods for biomolecular systems. *Angew. Chem. Int. Ed.* **48**(7), 1198–1229 (2009)
13. Tomasi, J., Persico, M.: Molecular interactions in solution – an overview of methods based on continuous distributions of the solvent. *Chem. Rev.* **94**(7), 2027–2094 (1994)
14. Tomasi, J., Mennucci, B., Laug, P.: The modeling and simulation of the liquid phase. In: Le Bris, C. (ed.) *Handbook of Numerical Analysis: Special Volume. Computational Chemistry*, pp. 271–323. Elsevier, Amsterdam (2003)
15. Tomasi, J., Mennucci, B., Cammi, R.: Quantum mechanical continuum solvation models. *Chem. Rev.* **105**(8), 2999–3093 (2005)

## Lobatto Methods

Laurent O. Jay  
 Department of Mathematics, The University of Iowa,  
 Iowa City, IA, USA

## Introduction

Lobatto methods for the numerical integration of differential equations are named after Reuel Lobatto. Reuel Lobatto (1796–1866) was a Dutch mathematician working most of his life as an advisor

for the government in the fields of life insurance and of weights and measures. In 1842, he was appointed professor of mathematics at the Royal Academy in Delft (known nowadays as Delft University of Technology). Lobatto methods are characterized by the use of approximations to the solution at the two end points  $t_n$  and  $t_{n+1}$  of each subinterval of integration  $[t_n, t_{n+1}]$ . Two well-known Lobatto methods based on the trapezoidal quadrature rule which are often used in practice are the *(implicit) trapezoidal rule* and the *Störmer-Verlet-leapfrog method*.

### The (Implicit) Trapezoidal Rule

Consider a system of ordinary differential equations (ODEs):

$$\frac{d}{dt}y = f(t, y) \quad (1)$$

where  $f : \mathbb{R} \times \mathbb{R}^d \rightarrow \mathbb{R}^d$ . Starting from  $y_0$  at  $t_0$  one step  $(t_n, y_n) \mapsto (t_{n+1}, y_{n+1})$  of the (implicit) trapezoidal rule applied to (1) is given by the implicit relation:

$$y_{n+1} = y_n + \frac{h_n}{2} (f(t_n, y_n) + f(t_{n+1}, y_{n+1}))$$

where  $h_n = t_{n+1} - t_n$  is the step size. The (implicit) trapezoidal rule is oftentimes called the *Crank-Nicholson method* when considered in the context of time-dependent partial differential equations (PDEs). This implicit method requires the solution of a system of  $d$  equations for  $y_{n+1} \in \mathbb{R}^d$  that can be expressed as:

$$F(y_{n+1}) := y_{n+1} - y_n - \frac{h_n}{2} (f(t_n, y_n) + f(t_{n+1}, y_{n+1})) = 0$$

and which is nonlinear when  $f(t, y)$  is nonlinear in  $y$ . Starting from an initial guess  $y_{n+1}^{(0)} \approx y_{n+1}$ , the solution  $y_{n+1}$  can be approximated iteratively by modified Newton iterations as follows:

$$y_{n+1}^{(k+1)} = y_{n+1}^{(k)} + p_{n+1}^{(k)}, \quad J_n p_{n+1}^{(k)} = -F(y_{n+1}^{(k)})$$

using, for example, an approximate Jacobian:

$$J_n = I_d - \frac{h_n}{2} D_y f(t_n, y_n) \approx D_y F(y_{n+1}^{(k)}).$$



Taking  $J_n = I_d$  leads to fixed-point iterations:

$$y_{n+1}^{(k+1)} = y_n + \frac{h_n}{2} \left( f(t_n, y_n) + f(t_{n+1}, y_{n+1}^{(k)}) \right).$$

### The Generalized Newton-Störmer-Verlet-Leapfrog Method

Consider now a partitioned system of ODEs:

$$\frac{d}{dt}q = v(t, p, q), \quad \frac{d}{dt}p = f(t, q, p) \quad (2)$$

where  $v : \mathbb{R} \times \mathbb{R}^{d_q} \times \mathbb{R}^{d_p} \rightarrow \mathbb{R}^{d_q}$  and  $f : \mathbb{R} \times \mathbb{R}^{d_q} \times \mathbb{R}^{d_p} \rightarrow \mathbb{R}^{d_p}$ . Starting from  $(q_0, p_0)$  at  $t_0$  one step  $(t_n, q_n, p_n) \mapsto (t_{n+1}, q_{n+1}, p_{n+1})$  of the *generalized Newton-Störmer-Verlet-leapfrog method* applied to (2) reads:

$$\begin{aligned} p_{n+1/2} &= p_n + \frac{h_n}{2} f(t_n, q_n, p_{n+1/2}), \\ q_{n+1} &= q_n + \frac{h_n}{2} \left( v(t_n, q_n, p_{n+1/2}) \right. \\ &\quad \left. + v(t_{n+1}, q_{n+1}, p_{n+1/2}) \right), \quad (3) \\ p_{n+1} &= p_{n+1/2} + \frac{h_n}{2} f(t_{n+1}, q_{n+1}, p_{n+1/2}) \end{aligned}$$

where  $h_n = t_{n+1} - t_n$  is the step size. The first equation is implicit for  $p_{n+1/2}$ , the second equation is implicit for  $q_{n+1}$ , and the last equation is explicit for  $p_{n+1}$ . When  $v(t, q, p) = v(t, p)$  is independent of  $q$ , and  $f(t, q, p) = f(t, q)$  is independent of  $p$  the method is fully explicit. If in addition  $v(t, q, p) = v(p)$  is independent of  $t$  and  $q$ , the method can be simply expressed as:

$$\begin{aligned} p_{n+1/2} &= p_n + \frac{h_n}{2} f(t_n, q_n), \\ q_{n+1} &= q_n + h_n v(p_{n+1/2}), \\ p_{n+1} &= p_{n+1/2} + \frac{h_n}{2} f(t_{n+1}, q_{n+1}). \end{aligned}$$

This explicit method is often applied as follows:

$$\begin{aligned} p_{n+1/2} &= p_{n-1/2} + \frac{1}{2}(h_{n-1} + h_n) f(t_n, q_n), \\ q_{n+1} &= q_n + h_n v(p_{n+1/2}). \end{aligned}$$

Depending on the field of applications, this method is known under different names: the *Störmer method* in astronomy; the *Verlet method* in molecular dynamics; the *leapfrog method* in the context of time-dependent PDEs, in particular for wave equations. This method can be traced back to Newton's Principia (1687), see [10].

### Lobatto Methods

In this entry, we consider families of Runge-Kutta (RK) methods based on Lobatto quadrature formulas whose simplest member is the trapezoidal quadrature rule. When applied to (1) Lobatto RK methods can be expressed as follows:

$$\begin{aligned} Y_{ni} &= y_n + h_n \sum_{j=1}^s a_{ij} f(t_n + c_j h, Y_{nj}) \\ &\text{for } i = 1, \dots, s, \quad (4) \end{aligned}$$

$$y_{n+1} = y_n + h_n \sum_{j=1}^s b_j f(t_n + c_j h, Y_{nj}) \quad (5)$$

where the stage value  $s$  satisfies  $s \geq 2$  and the coefficients  $a_{ij}, b_j, c_j$  characterize the Lobatto RK method. The  $s$  intermediate values  $Y_{nj}$  for  $j = 1, \dots, s$  are called the *internal stages* and can be considered as approximations to the solution at  $t_n + c_j h_n$ , the main numerical RK approximation at  $t_{n+1} = t_n + h_n$  is given by  $y_{n+1}$ . Lobatto RK methods are characterized by  $c_1 = 0$  and  $c_s = 1$ . They can also be considered in combination with other families of RK methods, for example, with Gauss methods in the context of certain systems of differential-algebraic equations (DAEs), see the section “[Lobatto Methods for DAEs](#)” below. The symbol III is usually found in the literature associated to Lobatto methods, the symbols I and II being reserved for the two types of Radau methods. The (implicit) trapezoidal rule is the simplest member ( $s = 2$ ) in the Lobatto IIIA family. The generalized Newton-Störmer-Verlet-leapfrog method seen above can be interpreted as a partitioned Runge-Kutta (PRK) resulting from the combination of the (implicit) trapezoidal rule and the Lobatto IIIB method for  $s = 2$ , see the section “[Additive Lobatto Methods for Split and Partitioned ODEs](#)” below.

### Families of Lobatto Methods

For a fixed value of  $s$ , the various families of Lobatto methods described below all share the same coefficients  $b_j, c_j$  of the corresponding Lobatto quadrature formula.

#### Lobatto Quadrature Formulas

The problem of approximating a Riemann integral:

$$\int_{t_n}^{t_n+h_n} f(t)dt \tag{6}$$

with  $f$  assumed to be continuous is equivalent to the problem of solving the initial value problem at  $t = t_n + h_n$ :

$$\frac{d}{dt}y = f(t), \quad y(t_n) = 0$$

since  $y(t_n + h_n) = \int_{t_n}^{t_n+h_n} f(t)dt$ . The integral (6) can be approximated by using a standard quadrature formula:

$$\int_{t_n}^{t_n+h_n} f(t)dt \approx h_n \left( \sum_{i=1}^s b_i f(t_n + c_i h_n) \right)$$

with  $s$  node coefficients  $c_1, \dots, c_s$ , and  $s$  weight coefficients  $b_1, \dots, b_s$ . Lobatto quadrature formulas, also known as Gauss-Lobatto quadrature formulas in the literature, are given for  $s \geq 2$  by a set of nodes and weights satisfying conditions described hereafter. The  $s$  nodes  $c_j$  are the roots of the polynomial of degree  $s$ :

$$\frac{d^{s-2}}{dt^{s-2}}(t^{s-1}(1-t)^{s-1}).$$

These nodes satisfy  $c_1 = 0 < c_2 < \dots < c_s = 1$ . The weights  $b_j$  and nodes  $c_j$  satisfy the condition  $B(2s-2)$  where:

$$B(p) : \sum_{j=1}^s b_j c_j^{k-1} = \frac{1}{k} \quad \text{for } k = 1, \dots, p,$$

implying that the quadrature formula is of order  $2s-2$ . There exists an explicit formula for the weights

$$b_j = \frac{1}{s(s-1)P_{s-1}(2c_j-1)^2} > 0$$

for  $j = 1, \dots, s$   $\left( b_1 = b_s = \frac{1}{s(s-1)} \right)$

where

$$P_k(x) = \frac{1}{k!2^k} \frac{d^k}{dx^k} ((x^2-1)^k)$$

is the  $k$ th Legendre polynomial. Lobatto quadrature formulas are symmetric, that is their nodes and weights satisfy:

$$b_{s+1-j} = b_j, \quad c_{s+1-j} = 1 - c_j \quad \text{for } j = 1, \dots, s.$$

For  $s = 3$ , we obtain the famous Simpson's rule:

$$(b_1, b_2, b_3) = (1/6, 2/3, 1/6), (c_1, c_2, c_3) = (0, 1/2, 1).$$

Procedures to compute numerically accurately the nodes and weights of high order Lobatto quadrature formulas can be found in [7] and [23]. The subroutine GQRUL from the IMSL/MATH-LIBRARY can compute numerically these nodes and weights.

#### Lobatto Families

The families of Lobatto RK methods differ only in the values of their coefficients  $a_{ij}$ . Various equivalent definitions can be found in the literature. The coefficients  $a_{ij}$  of these families can be linearly implicitly defined with the help of so-called *simplifying assumptions*:

$$C(q) : \sum_{j=1}^s a_{ij} c_j^{k-1} = \frac{c_i^k}{k}$$

for  $i = 1, \dots, s$  and  $k = 1, \dots, q,$

$$D(r) : \sum_{i=1}^s b_i c_i^{k-1} a_{ij} = \frac{b_j}{k} (1 - c_j^k)$$

for  $j = 1, \dots, s$  and  $k = 1, \dots, r.$

The importance of these simplifying assumptions comes from a fundamental result due to Butcher, see [5, 9], saying that a RK method satisfying the simplifying assumptions  $B(p)$ ,  $C(q)$ , and  $D(r)$  is of order at least  $\min(p, 2q+2, q+r+1)$ . The coefficients  $a_{ij}, b_j, c_j$  characterizing the Lobatto RK method (4) and (5) will be displayed below in the form of a table called a *Butcher-tableau*:



$$\begin{array}{c|cccccc}
c_1 = 0 & a_{11} & a_{12} & \cdots & a_{1,s-1} & a_{1s} \\
c_2 & a_{21} & a_{22} & \cdots & a_{2,s-1} & a_{2s} \\
\vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\
c_{s-1} & a_{s-1,1} & a_{s-1,2} & \cdots & a_{s-1,s-1} & a_{s-1,s} \\
c_s = 1 & a_{s1} & a_{s2} & \cdots & a_{s,s-1} & a_{ss} \\
\hline
& b_1 & b_2 & \cdots & b_{s-1} & b_s
\end{array}$$

In the four main families of Lobatto methods described below, namely Lobatto IIIA, Lobatto IIIB, Lobatto IIIC, and Lobatto IIIC\*, only one method does not satisfy the relation  $C(1)$ , that is,

$$\sum_{j=1}^s a_{ij} = c_i \quad \text{for } i = 1, \dots, s,$$

this is the Lobatto IIIB method for  $s = 2$ , see below. The Lobatto IIIA, IIIB, IIIC, and IIIC\* methods can all be interpreted as perturbed collocation methods [19] and discontinuous collocation methods [11].

#### Lobatto IIIA

The coefficients  $a_{ij}^A$  of Lobatto IIIA methods can be defined by  $C(s)$  (Table 1). They satisfy  $D(s-2)$ ,  $a_{sj}^A = b_j$  for  $j = 1, \dots, s$ , and  $a_{ij}^A = 0$  for  $j = 1, \dots, s$ . Lobatto IIIA methods are symmetric and of nonstiff order  $2s - 2$ . Their stability function  $R(z)$  is given by the  $(s-1, s-1)$ -Padé approximation to  $e^z$ . They are  $A$ -stable, but not  $L$ -stable since  $R(\infty) = (-1)^{s+1}$ . They are not  $B$ -stable and thus not algebraically stable. They can be interpreted as collocation methods. Since the first internal stage  $Y_{n1}$  of Lobatto IIIA methods is explicit ( $Y_{n1} = y_n$  and  $f(t_n + c_1 h_n, Y_{n1}) = f(t_n, y_n)$ ) and the last internal stage satisfies  $Y_{ns} = y_{n+1}$  (and thus  $f(t_{n+1}, y_{n+1}) = f(t_n + c_s h_n, Y_{ns})$ ), these methods are comparable in terms of computational work to Gauss methods with  $s - 1$  internal stages since they also have the same nonstiff order  $2s - 2$ . For  $s = 2$ , we obtain the (implicit) trapezoidal rule which is often expressed without its two internal stages  $Y_{n1}, Y_{n2}$  since they are respectively equal to  $y_n$  and  $y_{n+1}$ . The method for  $s = 3$  is sometimes called the *Hermite-Simpson (or Clippinger-Dimsdale) method* and it has been used, for example, in trajectory optimization problems [4]. This method can be equivalently expressed in a compact form as:

$$\begin{aligned}
Y_{n2} &= \frac{1}{2}(y_n + y_{n+1}) \\
&\quad + \frac{h_n}{8}(f(t_n, y_n) - f(t_{n+1}, y_{n+1})), \\
y_{n+1} &= y_n + \frac{h_n}{6}(f(t_n, y_n) + 4f(t_{n+1/2}, Y_{n2}) \\
&\quad + f(t_{n+1}, y_{n+1}))
\end{aligned}$$

where  $t_{n+1/2} = t_n + h_n/2$ . It can be even further reduced by rewriting

$$\begin{aligned}
y_{n+1} &= y_n + \frac{h_n}{6}(f(t_n, y_n) + f(t_{n+1}, y_{n+1})) \\
&\quad + \frac{2h_n}{3}f\left(t_{n+1/2}, \frac{1}{2}(y_n + y_{n+1})\right. \\
&\quad \left. + \frac{h_n}{8}(f(t_n, y_n) - f(t_{n+1}, y_{n+1}))\right).
\end{aligned}$$

#### Lobatto IIIB

The coefficients  $a_{ij}^B$  of Lobatto IIIB methods can be defined by  $D(s)$  (Table 2). They satisfy  $C(s-2)$ ,  $a_{i1}^B = b_1$  for  $i = 1, \dots, s$  and  $a_{is}^B = 0$  for  $i = 1, \dots, s$ . Lobatto IIIB methods are symmetric and of nonstiff order  $2s - 2$ . Their stability function  $R(z)$  is given by the  $(s-1, s-1)$ -Padé approximation to  $e^z$ . They are  $A$ -stable, but not  $L$ -stable since  $R(\infty) = (-1)^{s+1}$ . They are not  $B$ -stable and thus not algebraically stable. The coefficients  $a_{ij}^B$  can also be obtained from the coefficients  $a_{ij}^A$  of Lobatto IIIA through the relations:

$$b_i a_{ij}^B + b_j a_{ji}^A - b_i b_j = 0 \quad \text{for } i, j = 1, \dots, s,$$

or

$$a_{ij}^B = b_j - a_{s+1-i, s+1-j}^A \quad \text{for } i, j = 1, \dots, s.$$

#### Lobatto IIIC

The coefficients  $a_{ij}^C$  of Lobatto IIIC methods can be defined by  $a_{i1}^C = b_1$  for  $i = 1, \dots, s$  and  $C(s-1)$  (Table 3). They satisfy  $D(s-1)$  and  $a_{sj}^C = b_j$  for  $j = 1, \dots, s$ . Lobatto IIIC methods are of nonstiff order  $2s - 2$ . They are not symmetric. Their stability function  $R(z)$  is given by the  $(s-2, s)$ -Padé approximation to  $e^z$ . They are  $L$ -stable. They are algebraically stable and thus  $B$ -stable. They are excellent methods for stiff problems.

**Lobatto Methods, Table 1** Coefficients of Lobatto IIIA for  $s = 2, 3, 4, 5$

			0	0	0	0	0
0	0	0	0	0	0	0	0
1	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{5}{24}$	$\frac{1}{3}$	$-\frac{1}{24}$	$\frac{1}{2} - \frac{\sqrt{5}}{10}$
$A_{s=2}$	$\frac{1}{2}$	$\frac{1}{2}$	1	$\frac{1}{6}$	$\frac{2}{3}$	$\frac{1}{6}$	$\frac{1}{2} + \frac{\sqrt{5}}{10}$
			$A_{s=3}$	$\frac{1}{6}$	$\frac{2}{3}$	$\frac{1}{6}$	1
							$A_{s=4}$
							$\frac{1}{12}$
							$\frac{5}{12}$
							$\frac{5}{12}$
							$\frac{1}{12}$

  

	0	0	0	0	0
$\frac{1}{2} - \frac{\sqrt{21}}{14}$	$\frac{119 + 3\sqrt{21}}{1960}$	$\frac{343 - 9\sqrt{21}}{2520}$	$\frac{392 - 96\sqrt{21}}{2205}$	$\frac{343 - 69\sqrt{21}}{2520}$	$\frac{-21 + 3\sqrt{21}}{1960}$
$\frac{1}{2}$	$\frac{13}{320}$	$\frac{392 + 105\sqrt{21}}{2880}$	$\frac{8}{45}$	$\frac{392 - 105\sqrt{21}}{2880}$	$\frac{3}{320}$
$\frac{1}{2} + \frac{\sqrt{21}}{14}$	$\frac{119 - 3\sqrt{21}}{1960}$	$\frac{343 + 69\sqrt{21}}{2520}$	$\frac{392 + 96\sqrt{21}}{2205}$	$\frac{343 + 9\sqrt{21}}{2520}$	$\frac{-21 - 3\sqrt{21}}{1960}$
1	$\frac{1}{20}$	$\frac{49}{180}$	$\frac{16}{45}$	$\frac{49}{180}$	$\frac{1}{20}$
$A_{s=5}$	$\frac{1}{20}$	$\frac{49}{180}$	$\frac{16}{45}$	$\frac{49}{180}$	$\frac{1}{20}$

**Lobatto IIIC\***

Lobatto IIIC\* are also known as Lobatto III methods [5], Butcher’s Lobatto methods [9], and Lobatto IIIC methods [22] in the literature. (The name Lobatto IIIC\* was suggested by Robert P.K. Chan in an e-mail correspondence with the author on June 13, 1995.) The coefficients  $a_{ij}^{C*}$  of Lobatto IIIC\* methods can be defined by  $a_{is}^{C*} = 0$  for  $i = 1, \dots, s$  and  $C(s - 1)$  (Table 4). They satisfy  $D(s - 1)$  and  $a_{1j}^{C*} = 0$  for  $j = 1, \dots, s$ . Lobatto IIIC\* methods are of nonstiff order  $2s - 2$ . They are not symmetric. Their stability function  $R(z)$  is given by the  $(s, s - 2)$ -Padé approximation to  $e^z$ . They are not  $A$ -stable. They are not  $B$ -stable and thus not algebraically stable. The Lobatto IIIC\* method for  $s = 2$  is sometimes called the *explicit trapezoidal rule*. The coefficients  $a_{ij}^{C*}$  can also be obtained from the coefficients  $a_{ij}^C$  of Lobatto IIIC through the relations:

$$b_i a_{ij}^{C*} + b_j a_{ji}^C - b_i b_j = 0 \quad \text{for } i, j = 1, \dots, s,$$

or

$$a_{ij}^{C*} = b_j - a_{s+1-i, s+1-j}^C \quad \text{for } i, j = 1, \dots, s.$$

**Other Families of Lobatto Methods**

Most Lobatto methods of interest found in the literature can be expressed as linear combinations of the four fundamental Lobatto IIIA, IIIB, IIIC, and IIIC\* methods. In fact, one can consider a very general family of methods with three real parameters  $(\alpha_A, \alpha_B, \alpha_C)$  by considering Lobatto coefficients of the form:

$$a_{ij}(\alpha_A, \alpha_B, \alpha_C) = \alpha_A a_{ij}^A + \alpha_B a_{ij}^B + \alpha_C a_{ij}^C + \alpha_{C*} a_{ij}^{C*} \tag{7}$$

where  $\alpha_{C*} = 1 - \alpha_A - \alpha_B - \alpha_C$ . For any choice of  $(\alpha_A, \alpha_B, \alpha_C)$  the corresponding Lobatto RK method is of nonstiff order  $2s - 2$  [13]. The Lobatto IIIS methods presented in [6] depend on a real parameter  $\sigma$ . They can be expressed as:

$$a_{ij}^S(\sigma) = (1 - \sigma)(a_{ij}^A + a_{ij}^B) + \left(\sigma - \frac{1}{2}\right)(a_{ij}^C + a_{ij}^{C*})$$

for  $i, j = 1, \dots, s$ ,

corresponding to  $\alpha_A = \alpha_B = 1 - \sigma$  and  $\alpha_C = \alpha_{C*} = \sigma - \frac{1}{2}$  in (7). These methods satisfy  $C(s - 2)$  and



**Lobatto Methods, Table 2** Coefficients of Lobatto IIIB for  $s = 2, 3, 4, 5$

$B_{s=2} \begin{array}{c c} 0 & \frac{1}{2} \ 0 \\ 1 & \frac{1}{2} \ 0 \\ \hline B_{s=2} & \frac{1}{2} \ \frac{1}{2} \end{array}$	$B_{s=3} \begin{array}{c ccc} 0 & \frac{1}{6} & -\frac{1}{6} & 0 \\ \frac{1}{2} & \frac{1}{6} & \frac{1}{3} & 0 \\ 1 & \frac{1}{6} & \frac{5}{6} & 0 \\ \hline B_{s=3} & \frac{1}{6} & \frac{2}{3} & \frac{1}{6} \end{array}$	$B_{s=4} \begin{array}{c ccc c} 0 & \frac{1}{12} & \frac{-1-\sqrt{5}}{24} & \frac{-1+\sqrt{5}}{24} & 0 \\ \frac{1}{2} - \frac{\sqrt{5}}{10} & \frac{1}{12} & \frac{25+\sqrt{5}}{120} & \frac{25-13\sqrt{5}}{120} & 0 \\ \frac{1}{2} + \frac{\sqrt{5}}{10} & \frac{1}{12} & \frac{25+13\sqrt{5}}{120} & \frac{25-\sqrt{5}}{120} & 0 \\ 1 & \frac{1}{12} & \frac{11-\sqrt{5}}{24} & \frac{11+\sqrt{5}}{24} & 0 \\ \hline B_{s=4} & \frac{1}{12} & \frac{5}{12} & \frac{5}{12} & \frac{1}{12} \end{array}$			
$B_{s=5} \begin{array}{c ccccc} 0 & \frac{1}{20} & \frac{-7-\sqrt{21}}{120} & \frac{1}{15} & \frac{-7+\sqrt{21}}{120} & 0 \\ \frac{1}{2} - \frac{\sqrt{21}}{14} & \frac{1}{20} & \frac{343+9\sqrt{21}}{2520} & \frac{56-15\sqrt{21}}{315} & \frac{343-69\sqrt{21}}{2520} & 0 \\ \frac{1}{2} & \frac{1}{20} & \frac{49+12\sqrt{21}}{360} & \frac{8}{45} & \frac{49-12\sqrt{21}}{360} & 0 \\ \frac{1}{2} + \frac{\sqrt{21}}{14} & \frac{1}{20} & \frac{343+69\sqrt{21}}{2520} & \frac{56+15\sqrt{21}}{315} & \frac{343-9\sqrt{21}}{2520} & 0 \\ 1 & \frac{1}{20} & \frac{119-3\sqrt{21}}{360} & \frac{13}{45} & \frac{119+3\sqrt{21}}{360} & 0 \\ \hline B_{s=5} & \frac{1}{20} & \frac{49}{180} & \frac{16}{45} & \frac{49}{180} & \frac{1}{20} \end{array}$					

$D(s - 2)$ . They are symmetric and symplectic. Their stability function  $R(z)$  is given by the  $(s - 1, s - 1)$ -Padé approximation to  $e^z$ . They are  $A$ -stable, but not  $L$ -stable. They are algebraically stable and thus  $B$ -stable. The Lobatto IIIS coefficients for  $\sigma = 1/2$  are given by:

$$a_{ij}^S(1/2) = \frac{1}{2} (a_{ij}^A + a_{ij}^B) \quad \text{for } i, j = 1, \dots, s.$$

For  $\sigma = 1$  we obtain the Lobatto IIID methods [6, 13]:

$$a_{ij}^D = a_{ij}^S(1) = \frac{1}{2} (a_{ij}^C + a_{ij}^{C*}) \quad \text{for } i, j = 1, \dots, s.$$

These methods are called Lobatto III $_E$  in [19] and Lobatto III $_E$  in [22]. They satisfy  $C(s-1)$  and  $D(s-1)$ , and they can be interpreted as perturbed collocation methods [19]. Another family of Lobatto RK methods is given by the Lobatto III $_D$  family of [19] called here Lobatto III $_D$  where the coefficients for  $s = 2, 3$  are given in Table 5. (Notice on p. 205 of [19] that  $\gamma_1 = -4(2m - 1)$ .) These methods correspond to

$\alpha_A = 2, \alpha_B = 2, \alpha_C = -1$ , and  $\alpha_{C^*} = -2$  in (7). Their stability function  $R(z)$  is given by the  $(s - 2, s)$ -Padé approximation to  $e^z$ . These methods are  $L$ -stable. They are algebraically stable and thus  $B$ -stable. They are of nonstiff order  $2s - 2$ . They are not symmetric. They can be interpreted as perturbed collocation methods [19].

### Additive Lobatto Methods for Split and Partitioned ODEs

Consider a split system of ODEs:

$$\frac{d}{dt}y = f_1(t, y) + f_2(t, y) \tag{8}$$

where  $f_1, f_2 : \mathbb{R} \times \mathbb{R}^d \rightarrow \mathbb{R}^d$ . Starting from  $y_0$  at  $t_0$  one step  $(t_n, y_n) \mapsto (t_{n+1}, y_{n+1})$  of an additive Lobatto RK method applied to (8) reads:

**Lobatto Methods, Table 3** Coefficients of Lobatto IIIC for  $s = 2, 3, 4, 5$

$0$	$\frac{1}{2} - \frac{1}{2}$	$0$	$\frac{1}{6} - \frac{1}{3} \quad \frac{1}{6}$	$0$	$\frac{1}{12} \quad -\frac{\sqrt{5}}{12} \quad \frac{\sqrt{5}}{12} \quad -\frac{1}{12}$
$1$	$\frac{1}{2} \quad \frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{6} \quad \frac{5}{12} \quad -\frac{1}{12}$	$\frac{1}{2} - \frac{\sqrt{5}}{10}$	$\frac{1}{12} \quad \frac{1}{4} \quad \frac{10 - 7\sqrt{5}}{60} \quad \frac{\sqrt{5}}{60}$
$C_{s=2}$	$\frac{1}{2} \quad \frac{1}{2}$	$1$	$\frac{1}{6} \quad \frac{2}{3} \quad \frac{1}{6}$	$\frac{1}{2} + \frac{\sqrt{5}}{10}$	$\frac{1}{12} \quad \frac{10 + 7\sqrt{5}}{60} \quad \frac{1}{4} \quad -\frac{\sqrt{5}}{60}$
		$C_{s=3}$	$\frac{1}{6} \quad \frac{2}{3} \quad \frac{1}{6}$	$1$	$\frac{1}{12} \quad \frac{5}{12} \quad \frac{5}{12} \quad \frac{1}{12}$
				$C_{s=4}$	$\frac{1}{12} \quad \frac{5}{12} \quad \frac{5}{12} \quad \frac{1}{12}$
$0$	$\frac{1}{20}$	$0$	$-\frac{7}{60}$	$\frac{2}{15}$	$-\frac{7}{60}$
$\frac{1}{2} - \frac{\sqrt{21}}{14}$	$\frac{1}{20}$	$\frac{1}{2}$	$\frac{29}{180}$	$\frac{47 - 15\sqrt{21}}{315}$	$\frac{203 - 30\sqrt{21}}{1260}$
$\frac{1}{2}$	$\frac{1}{20}$	$\frac{1}{2} + \frac{\sqrt{21}}{14}$	$\frac{329 + 105\sqrt{21}}{2880}$	$\frac{73}{360}$	$\frac{329 - 105\sqrt{21}}{2880}$
$\frac{1}{2} + \frac{\sqrt{21}}{14}$	$\frac{1}{20}$	$1$	$\frac{203 + 30\sqrt{21}}{1260}$	$\frac{47 + 15\sqrt{21}}{315}$	$\frac{29}{180}$
$1$	$\frac{1}{20}$	$C_{s=5}$	$\frac{49}{180}$	$\frac{16}{45}$	$\frac{49}{180}$
					$\frac{1}{20}$

$$Y_{ni} = y_n + h_n \sum_{j=1}^s (a_{1,ij} f_1(t_n + c_j h, Y_{nj}) + a_{2,ij} f_2(t_n + c_j h, Y_{nj}))$$

for  $i = 1, \dots, s$ ,

$$y_{n+1} = y_n + h_n \sum_{j=1}^s b_j (f_1(t_n + c_j h, Y_{nj}) + f_2(t_n + c_j h, Y_{nj}))$$

where  $s \geq 2$  and the coefficients  $a_{1,ij}, a_{2,ij}, b_j, c_j$  characterize the additive Lobatto RK method. Consider, for example, any coefficients  $a_{1,ij}$  and  $a_{2,ij}$  from the family (7), the additive method is of nonstiff order  $2s - 2$  [13]. The partitioned system of ODEs (2) can be expressed in the form (8) by having  $d = d_q + d_p$ ,  $y = (q, p) \in \mathbb{R}^{d_q} \times \mathbb{R}^{d_p}$ , and:

$$f_1(t, q, p) = \begin{pmatrix} v(t, q, p) \\ 0 \end{pmatrix},$$

$$f_2(t, q, p) = \begin{pmatrix} 0 \\ f(t, q, p) \end{pmatrix}.$$

Applying for  $s = 2$  the Lobatto IIIA coefficients as  $a_{1,ij}$  and the Lobatto IIIB coefficients as  $a_{2,ij}$ , we obtain again the generalized Newton-Störmer-Verlet-leapfrog method (3). Additive Lobatto methods have been considered in multibody dynamics in [13, 21]. Additive methods are more general than partitioned methods since partitioned system of ODEs can always be reformulated as a split system of ODEs, but the reverse is false in general.

### Lobatto Methods for DAEs

An important use of Lobatto methods is for the solution of differential-algebraic equations (DAEs). DAEs consist generally of coupled systems of differential equations and nonlinear relations. They arise typically in mechanics and electrical/electronic circuits simulation.





**Lobatto Methods, Table 4** Coefficients of Lobatto IIIC\* for  $s = 2, 3, 4, 5$

			0	0	0	0	0		
0	0 0	0	0 0 0	$\frac{1}{2} - \frac{\sqrt{5}}{10}$	$\frac{5 + \sqrt{5}}{60}$	$\frac{1}{6}$	$\frac{15 - 7\sqrt{5}}{60}$	0	
1	1 0	$\frac{1}{2}$	$\frac{1}{4} \frac{1}{4} 0$	$\frac{1}{2} + \frac{\sqrt{5}}{10}$	$\frac{5 - \sqrt{5}}{60}$	$\frac{15 + 7\sqrt{5}}{60}$	$\frac{1}{6}$	0	
$C_{s=2}^*$	$\frac{1}{2} \frac{1}{2}$	1	0 1 0	1	$\frac{1}{6}$	$\frac{5 - \sqrt{5}}{12}$	$\frac{5 + \sqrt{5}}{12}$	0	
		$C_{s=3}^*$	$\frac{1}{6} \frac{2}{3} \frac{1}{6}$		$C_{s=4}^*$	$\frac{1}{12}$	$\frac{5}{12}$	$\frac{5}{12}$	$\frac{1}{12}$
	0	0	0	0	0	0	0	0	
	$\frac{1}{2} - \frac{\sqrt{21}}{14}$	$\frac{1}{14}$	$\frac{1}{9}$	$\frac{13 - 3\sqrt{21}}{63}$	$\frac{14 - 3\sqrt{21}}{126}$			0	
	$\frac{1}{2}$	$\frac{1}{32}$	$\frac{91 + 21\sqrt{21}}{576}$	$\frac{11}{72}$	$\frac{91 - 21\sqrt{21}}{576}$			0	
	$\frac{1}{2} + \frac{\sqrt{21}}{14}$	$\frac{1}{14}$	$\frac{14 + 3\sqrt{21}}{126}$	$\frac{13 + 3\sqrt{21}}{63}$	$\frac{1}{9}$			0	
	1	0	$\frac{7}{18}$	$\frac{2}{9}$	$\frac{7}{18}$			0	
$C_{s=5}^*$	$\frac{1}{20}$	$\frac{49}{180}$	$\frac{16}{45}$	$\frac{49}{180}$	$\frac{1}{20}$				

**Lobatto Methods, Table 5** Coefficients of Lobatto IIINW for  $s = 2, 3$  [19]

		0	$\frac{1}{6}$	0	$-\frac{1}{6}$
0	$\frac{1}{2} \frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{12}$	$\frac{5}{12}$	0
1	$-\frac{1}{2} \frac{1}{2}$	1	$\frac{1}{2}$	$\frac{1}{3}$	$\frac{1}{6}$
	$\frac{1}{2} \frac{1}{2}$		$\frac{1}{6}$	$\frac{2}{3}$	$\frac{1}{6}$

$$\begin{aligned}
 Y_{n1} &= y_n + \frac{h_n}{4}(f(t_n, Y_{n1}, \Lambda_{n1}) - f(t_{n+1}, Y_{n2}, \Lambda_{n2})), \\
 Y_{n2} &= y_n + \frac{h_n}{4}(3f(t_n, Y_{n1}, \Lambda_{n1}) + f(t_{n+1}, Y_{n2}, \Lambda_{n2})), \\
 y_{n+1} &= y_n + \frac{h_n}{2}(f(t_n, Y_{n1}, \Lambda_{n1}) + f(t_{n+1}, Y_{n2}, \Lambda_{n2})), \\
 0 &= \frac{1}{2}(k(t_n, Y_{n1}) + k(t_{n+1}, Y_{n2})), \\
 0 &= k(t_{n+1}, y_{n+1}).
 \end{aligned}$$

Consider, for example, a system of DAEs of the form:

$$\frac{d}{dt}y = f(t, y, \lambda), \quad 0 = k(t, y)$$

where  $D_y k(t, y) D_\lambda f(t, y, \lambda)$  is nonsingular. Lobatto methods can be applied to this class of problems while preserving their classical order of convergence [14]. For example, the application of the two-stage Lobatto IIID method can be expressed as:

For such DAEs, a combination of Gauss and Lobatto coefficients is also considered in [18]. Consider now overdetermined system of DAEs (ODAEs) of the form:

$$\begin{aligned}
 \frac{d}{dt}q &= v(t, q, p), \quad \frac{d}{dt}p = f(t, q, p, \lambda), \quad 0 = g(t, q), \\
 0 &= D_t g(t, q) + D_q g(t, q) v(t, q, p)
 \end{aligned} \tag{9}$$

where  $D_q g(t, q) D_p v(t, q, p) D_\lambda f(t, q, p, \lambda)$  is nonsingular. Very general Lobatto methods can be applied to this type of ODAEs [13]. Hamiltonian and



Lagrangian systems with holonomic constraints can be expressed in the form (9). For such ODAEs, the application of Lobatto IIIA and IIIB methods can be shown to preserve their classical order of convergence, to be variational integrators, and to preserve a symplectic two-form [8, 11, 12, 17]. For example, the application of the two-stage Lobatto IIIA and IIIB method reads:

$$\begin{aligned} q_{n+1} &= q_n + \frac{h_n}{2} \left( v(t_n, q_n, p_{n+1/2}) \right. \\ &\quad \left. + v(t_{n+1}, q_{n+1}, p_{n+1/2}) \right), \\ p_{n+1/2} &= p_n + \frac{h_n}{2} f(t_n, q_n, p_{n+1/2}, \Lambda_{n1}), \\ 0 &= g(t_{n+1}, q_{n+1}), \\ p_{n+1} &= p_{n+1/2} + \frac{h_n}{2} f(t_{n+1}, q_{n+1}, p_{n+1/2}, \Lambda_{n2}) \\ 0 &= D_t g(t_{n+1}, q_{n+1}) \\ &\quad + D_q g(t_{n+1}, q_{n+1}) v(t_{n+1}, q_{n+1}, p_{n+1}). \end{aligned}$$

Gauss methods with  $s$  stages can also be applied in combination with Lobatto methods with  $s+1$  stages for this type of ODAEs when  $f(t, q, p, \lambda)$  is decomposed in  $f(t, q, p) + r(t, q, \lambda)$  and they also possess these aforementioned properties while generally requiring less computational effort [15]. For example, the application of the midpoint-trapezoidal method (the (1, 1)-Gauss-Lobatto SPARK method of Jay [15]) reads:

$$\begin{aligned} Q_{n1} &= q_n + \frac{h_n}{2} v(t_{n+1/2}, Q_{n1}, P_{n1}) = \frac{1}{2}(q_n + q_{n+1}), \\ P_{n1} &= p_n + \frac{h_n}{2} f(t_{n+1/2}, Q_{n1}, P_{n1}) \\ &\quad + \frac{h_n}{2} r(t_n, q_n, \Lambda_{n1}), \\ q_{n+1} &= q_n + h_n v(t_{n+1/2}, Q_{n1}, P_{n1}), \\ p_{n+1} &= p_n + h_n f(t_{n+1/2}, Q_{n1}, P_{n1}) \\ &\quad + h_n \left( \frac{1}{2} r(t_n, q_n, \Lambda_{n1}) + \frac{1}{2} r(t_{n+1}, q_{n+1}, \Lambda_{n2}) \right), \\ 0 &= g(t_{n+1}, q_{n+1}), \\ 0 &= D_t g(t_{n+1}, q_{n+1}) \\ &\quad + D_q g(t_{n+1}, q_{n+1}) v(t_{n+1}, q_{n+1}, p_{n+1}). \end{aligned}$$

## Lobatto Methods for Some Other Classes of Problems

Lobatto IIIA methods have been considered for boundary value problems (BVP) due to their good stability properties [1, 2]. The *MATLAB* code `bvp4c` for BVP is based on three-stage collocation at Lobatto points, hence it is equivalent to the three-stage Lobatto IIIA method [16]. Lobatto methods have also been applied to delay differential equations (DDEs) [3]. The combination of Lobatto IIIA and IIIB methods has also been considered for the discrete multisymplectic integration of certain Hamiltonian partial differential equations (PDEs) such as the nonlinear Schrödinger equation and certain nonlinear wave equations [20].

## References

1. Ascher, U.M., Mattheij, R.M.M., Russell, R.D.: Numerical Solution of Boundary Value Problems for Ordinary Differential Equations. Classics in Applied Mathematics, vol. 13. SIAM, Philadelphia (1995)
2. Bashir-Ali, Z., Cash, J.R., Silva, H.H.M.: Lobatto deferred correction for stiff two-point boundary value problems. *Comput. Math. Appl.* **36**, 59–69 (1998)
3. Bellen, A., Guglielmi, N., Ruehli, A.E.: Methods for linear systems of circuit delay differential equations of neutral type. *IEEE Trans. Circuits Syst.* **46**, 212–216 (1999)
4. Betts, J.T.: Practical Methods for Optimal Control and Estimation Using Nonlinear Programming. Advances in Design and Control, 2nd edn. SIAM, Philadelphia (2008)
5. Butcher, J.C.: Numerical Methods for Ordinary Differential Equations, 2nd edn. Wiley, Chichester (2008)
6. Chan, R.P.K.: On symmetric Runge-Kutta methods of high order. *Computing* **45**, 301–309 (1990)
7. Gautschi, W.: High-order Gauss-Lobatto formulae. *Numer. Algorithms* **25**, 213–222 (2000)
8. Hairer, E., Wanner, G.: Solving Ordinary Differential Equations II. Stiff and Differential-Algebraic Problems. Computational Mathematics, vol. 14, 2nd edn. Springer, Berlin (1996)
9. Hairer, E., Nørsett, S.P., Wanner, G.: Solving Ordinary Differential Equations I. Nonstiff Problems. Computational Mathematics, vol. 18, 2nd edn. Springer, Berlin (1993)
10. Hairer, E., Lubich, C., Wanner, G.: Geometric numerical integration illustrated by the Störmer/Verlet method. *Acta Numer.* vol. 12, 1–51 (2003)
11. Hairer, E., Lubich, C., Wanner, G.: Geometric Numerical Integration. Computational Mathematics, vol. 31, 2nd edn. Springer, Berlin (2006)
12. Jay, L.O.: Symplectic partitioned Runge-Kutta methods for constrained Hamiltonian systems. *SIAM J. Numer. Anal.* **33**, 368–387 (1996)
13. Jay, L.O.: Structure preservation for constrained dynamics with super partitioned additive Runge-Kutta methods. *SIAM J. Sci. Comput.* **20**, 416–446 (1998)

14. Jay, L.O.: Solution of index 2 implicit differential-algebraic equations by Lobatto Runge-Kutta methods. *BIT* **43**, 91–104 (2003)
15. Jay, L.O.: Specialized partitioned additive Runge-Kutta methods for systems of overdetermined DAEs with holonomic constraints. *SIAM J. Numer. Anal.* **45**, 1814–1842 (2007)
16. Kierzenka, J., Shampine, L.F.: A BVP solver based on residual control and the MATLAB PSE. *ACM Trans. Math. Softw.* **27**, 299–316 (2001)
17. Leimkuhler, B., Reich, S.: *Simulating Hamiltonian Dynamics*, Cambridge Monographs on Applied and Computational Mathematics, vol. 14. Cambridge University Press, Cambridge (2005)
18. Murua, A.: Partitioned Runge-Kutta methods for semi-explicit differential-algebraic systems of index 2. Technical Report EHU-KZAA-IKT-196, University of the Basque country (1996)
19. Nørsett, S.P., Wanner, G.: Perturbed collocation and Runge-Kutta methods. *Numer. Math.* **38**, 193–208 (1981)
20. Ryland, B.N., McLachlan, R.I.: On multisymplecticity of partitioned Runge-Kutta methods. *SIAM J. Sci. Comput.* **30**, 1318–1340 (2008)
21. Schaub, M., Simeon, B.: Blended Lobatto methods in multi-body dynamics. *Z Angew. Math. Mech.* **83**, 720–728 (2003)
22. Sun, G.: A simple way of constructing symplectic Runge-Kutta methods. *J. Comput. Math.* **18**, 61–68 (2000)
23. von Matt, U.: Gauss quadrature. In: Gander, W., Hřebíček, J. (eds.) *Solving Problems in Scientific Computing Using Maple and Matlab*, vol. 14, 4th edn. Springer, Berlin (2004)

as well as boundary value problems and their discretizations. Some special fields in mathematics, such as semigroup theory, rely on notions that are strongly related to the logarithmic norm.

Let  $|\cdot|$  denote an arbitrary vector norm on  $\mathbb{C}^d$ , as well as its subordinate operator norm on  $\mathbb{C}^{d \times d}$ . The classical definition of the *logarithmic norm* of  $A \in \mathbb{C}^{d \times d}$  is

$$M[A] = \lim_{h \rightarrow 0^+} \frac{|I + hA| - 1}{h}. \quad (1)$$

It is easily computed for the most common norms, see Table 1. In Hilbert space, where the norm is generated by an inner product  $|x|^2 = \langle x, x \rangle$ , one may alternatively define the *least upper bound* logarithmic norm  $M[A]$  and the *greatest lower bound* logarithmic norm  $m[A]$  such that for all  $x$

$$m[A] \cdot |x|^2 \leq \operatorname{Re} \langle x, Ax \rangle \leq M[A] \cdot |x|^2. \quad (2)$$

Unlike (1), this also admits *unbounded operators*, while still agreeing with (1) if  $A$  is bounded, in which case it also holds that

$$m[A] = \lim_{h \rightarrow 0^-} \frac{|I + hA| - 1}{h}. \quad (3)$$

The functionals  $M[\cdot]$  and  $m[\cdot]$  can further be extended to nonlinear maps, both in a Banach and a Hilbert space setting, so that the above definitions become special cases for linear operators.

The logarithmic norm has a large number of useful properties and satisfy several important inequalities. For  $A, B \in \mathbb{C}^{d \times d}$ ,  $\alpha \in \mathbb{R}$  and  $z \in \mathbb{C}$ , some of the most important are:

1.  $-\operatorname{glb}[A] \leq M[A] \leq |A|$
2.  $M[\alpha A] = \alpha M[A]$ ,  $\alpha \geq 0$
3.  $M[A + zI] = M[A] + \operatorname{Re} z$
4.  $m[A] = -M[-A]$
5.  $M[A] + m[B] \leq M[A + B] \leq M[A] + M[B]$
6.  $|M[A] - M[B]| \leq |A - B|$
7.  $|m[A] - m[B]| \leq |A - B|$
8.  $e^{tm[A]} \leq |e^{tA}| \leq e^{tM[A]}$ ,  $t \geq 0$
9.  $M[A] < 0 \Rightarrow |A^{-1}| \leq -1/M[A]$
10.  $m[A] > 0 \Rightarrow |A^{-1}| \leq 1/m[A]$ .

## Logarithmic Norms

Gustaf Söderlind

Centre for Mathematical Sciences, Numerical Analysis, Lund University, Lund, Sweden

## Introduction

The *logarithmic norm* is a real-valued functional on operators, quantifying the notions of *definiteness* for matrices; *monotonicity* for nonlinear maps; and *ellipticity* for differential operators. It is defined either in terms of an inner product in Hilbert space, or in terms of the operator norm on a Banach space.

The logarithmic norm has a wide range of applications in matrix theory, stability theory, and numerical analysis. It offers various quantitative bounds on (functions of) operators, operator spectra, resolvents, Rayleigh quotients, and the numerical range. It also offers error bounds and stability estimates in initial