SELBERG'S ORTHOGONALITY CONJECTURE FOR AUTOMORPHIC L -FUNCTIONS

JIAN YA LIU¹ AND YANG BO YE²

ABSTRACT. Let π and π' be automorphic irreducible unitary cuspidal representations of $GL_m(\mathbb{Q}_{\mathbb{A}})$ and $GL_{m'}(\mathbb{Q}_{\mathbb{A}})$, respectively. Assume that π and π' are self contragredient. Under the Ramanujan conjecture on π and π' , we deduce a prime number theorem for $L(s, \pi \times \tilde{\pi}')$, which can be used to asymptotically describe whether $\pi' \cong \pi$, or $\pi' \cong \pi \otimes |\det(\cdot)|^{i\tau_0}$ for some non-zero $\tau_0 \in \mathbb{R}$, or $\pi' \not\cong \pi \otimes |\det(\cdot)|^{it}$ for any $t \in \mathbb{R}$. As a consequence, we prove the Selberg orthogonality conjecture, in a more precise form, for automorphic L -functions $L(s, \pi)$ and $L(s, \pi')$, under the Ramanujan conjecture. When $m = m' = 2$ and π and π' are representations corresponding to holomorphic cusp forms, our results are unconditional.

1. Introduction. Let π be an irreducible unitary cuspidal representation of $GL_m(\mathbb{Q}_{\mathbb{A}})$. Then the global L -function attached to π is given by products of local factors for $\text{Re } s > 1$ (Godement and Jacquet [3]):

$$L(s, \pi) = \prod_p L_p(s, \pi_p),$$

$$\Phi(s, \pi) = L_{\infty}(s, \pi_{\infty})L(s, \pi),$$

where

$$L_p(s, \pi_p) = \prod_{j=1}^m (1 - \alpha_{\pi}(p, j)p^{-s})^{-1}$$

and

$$L_{\infty}(s, \pi_{\infty}) = \prod_{j=1}^m \Gamma_{\mathbb{R}}(s + \mu_{\pi}(j)).$$

Here $\Gamma_{\mathbb{R}}(s) = \pi^{-s/2}\Gamma(s/2)$, and $\alpha_{\pi}(p, j)$ and $\mu_{\pi}(j)$, $j = 1, \dots, m$, are complex numbers associated with π_p and π_{∞} , respectively, according to the Langlands correspondence. De-

2000 Mathematics Subject Classification (MSC) 11F70, 11M26, 11M41.

¹Partially supported by China National Natural Science Foundation grant # 10125101.

²Project sponsored by the USA National Security Agency under Grant Number MDA904-03-1-0066. The United States Government is authorized to reproduce and distribute reprints notwithstanding any copyright notation herein.

note

$$(1.1) \quad a_\pi(p^k) = \sum_{1 \leq j \leq m} \alpha_\pi(p, j)^k.$$

Then for $\text{Re } s > 1$, we have

$$(1.2) \quad \frac{d}{ds} \log L(s, \pi) = - \sum_{n \geq 1} \frac{\Lambda(n) a_\pi(n)}{n^s},$$

where $\Lambda(n) = \log p$ if $n = p^k$ and $= 0$ otherwise. If π' is an automorphic irreducible cuspidal representation of $GL_{m'}(\mathbb{Q}_\mathbb{A})$, we define $L(s, \pi')$, $\alpha_{\pi'}(p, i)$, $\mu_{\pi'}(i)$, and $a_{\pi'}(p^k)$ likewise, for $i = 1, \dots, m'$. If π and π' are equivalent, then $m = m'$ and $\{\alpha_\pi(p, j)\} = \{\alpha_{\pi'}(p, i)\}$ for any p . Hence $a_\pi(n) = a_{\pi'}(n)$ for any $n = p^k$, when $\pi \cong \pi'$.

The Selberg orthogonality conjecture for automorphic L -functions $L(s, \pi)$ was proposed in 1989 (Selberg [19]). See also Ram Murty [15] [16].

CONJECTURE 1.1. (i) *For any automorphic irreducible cuspidal representation π of $GL_m(\mathbb{Q}_\mathbb{A})$*

$$(1.3) \quad \sum_{p \leq x} \frac{|a_\pi(p)|^2}{p} = \log \log x + O(1).$$

(ii) *For any automorphic irreducible cuspidal representations π and π' of $GL_m(\mathbb{Q}_\mathbb{A})$ and $GL_{m'}(\mathbb{Q}_\mathbb{A})$, respectively,*

$$(1.4) \quad \sum_{p \leq x} \frac{a_\pi(p) \bar{a}_{\pi'}(p)}{p} \ll 1,$$

if π is not equivalent to π' .

The asymptotic formula in (1.3) was proved by Rudnick and Sarnak [17] under a conjecture on the convergence of a series on prime powers (Hypothesis H below), and unconditionally for $m \leq 4$.

HYPOTHESIS H. *For $k \geq 2$,*

$$\sum_p \frac{|a_\pi(p^k)|^2 \log^2 p}{p^k} < \infty.$$

This Hypothesis H is trivial for $m = 1$, and follows from bounds toward the Ramanujan conjecture for $m = 2$. For $m = 3$ it was proved by Rudnick and Sarnak [17], while the case of $m = 4$ was proved by Kim and Sarnak [9]. For $m > 4$, Hypothesis H is an easy consequence of the Ramanujan conjecture. In this paper, we will assume the Ramanujan conjecture for primes p :

CONJECTURE 1.2. *Let π be an irreducible unitary cuspidal representation of $GL_m(\mathbb{Q}_\mathbb{A})$. For any unramified p , we have*

$$(1.5) \quad |\alpha_\pi(p, j)| = 1.$$

Note that in Conjecture 1.2 we do not include the Archimedean Ramanujan conjecture, $\operatorname{Re} \mu_\pi(j) = 0$.

What we will prove as a consequence of Conjecture 1.2 is the following orthogonality. Denote $\alpha(g) = |\det(g)|$.

THEOREM 1.3. *Let π and π' be irreducible unitary cuspidal representations of $GL_m(\mathbb{Q}_\mathbb{A})$ and $GL_{m'}(\mathbb{Q}_\mathbb{A})$, respectively, such that at least one of π and π' is self contragredient: $\pi \cong \tilde{\pi}$ or $\pi' \cong \tilde{\pi}'$. Assume the Ramanujan Conjecture 1.2 for both π and π' . Then*

$$\begin{aligned} \sum_{n \leq x} (\log n) \Lambda(n) a_\pi(n) \bar{a}_{\pi'}(n) &= \frac{x^{1+i\tau_0}}{1+i\tau_0} \log x - \frac{x^{1+i\tau_0}}{(1+i\tau_0)^2} + O\{x \exp(-c\sqrt{\log x})\} \\ &\quad \text{if } \pi' \cong \pi \otimes \alpha^{i\tau_0} \text{ for some } \tau_0 \in \mathbb{R}; \\ &= O\{x \exp(-c\sqrt{\log x})\} \\ &\quad \text{if } \pi' \not\cong \pi \otimes \alpha^{it} \text{ for any } t \in \mathbb{R}, \end{aligned}$$

where c is a positive constant.

Note that for $m = m' = 2$ and π and π' being representations corresponding to holomorphic cusp forms, i.e., when their Archimedean local components are discrete series or limits of discrete series, the Ramanujan conjecture was proved by Deligne. Therefore in this case, Theorem 1.3 is an unconditional result.

Theorem 1.3 is indeed a version of prime number theorem for the Rankin-Selberg L -function $L(s, \pi \times \pi')$. It is stronger than the following Mertens theorem on orthogonality with a weight $(\log n) \Lambda(n)/n$:

COROLLARY 1.4. *Let π and π' be given as in Theorem 1.3. Assume either (i) Ramanujan Conjecture 1.2 for both π and π' , or (ii) that $m = m' = 2$ and π and π' are*

representations corresponding to holomorphic cusp forms. Then

$$\begin{aligned}
(1.6) \quad \sum_{n \leq x} \frac{(\log n) \Lambda(n) a_\pi(n) \bar{a}_{\pi'}(n)}{n} &= \frac{1}{2} \log^2 x + c_1 + O\{\exp(-c\sqrt{\log x})\} \\
&\text{if } \pi' \cong \pi; \\
&= \frac{x^{i\tau_0}}{i\tau_0} \log x + \frac{x^{i\tau_0} - 1}{\tau_0^2} + c_2 + O\{\exp(-c\sqrt{\log x})\} \\
&\text{if } \pi' \cong \pi \otimes \alpha^{i\tau_0} \text{ for some } \tau_0 \in \mathbb{R}^\times; \\
&= c_3 + O\{\exp(-c\sqrt{\log x})\} \\
&\text{if } \pi' \not\cong \pi \otimes \alpha^{it} \text{ for any } t \in \mathbb{R},
\end{aligned}$$

where $c, c_1, \dots, c_3$ are positive constants.

A remarkable feature of this corollary is that it describes the orthogonality of $a_\pi(n)$ and $a_{\pi'}(n)$ in three cases with different main terms. As we are assuming Ramanujan and hence the Hypothesis H, we can control sums over prime powers and easily get an orthogonality over primes. Selberg's orthogonality conjecture 1.1 is then a consequence of Corollary 1.4 by partial summation.

COROLLARY 1.5 (SELBERG'S ORTHOGONALITY). *Let π and π' be given as in Theorem 1.3. Assume either (i) Ramanujan Conjecture 1.2 for both π and π' , or (ii) $m = m' = 2$ and that π and π' are representations corresponding to holomorphic cusp forms. Then*

$$\begin{aligned}
(1.7) \quad \sum_{p \leq x} \frac{a_\pi(p) \bar{a}_{\pi'}(p)}{p} &= \log \log x + c_4 + O\{\exp(-c\sqrt{\log x})\} \\
&\text{if } \pi' \cong \pi; \\
&= c_5 + \text{Ei}(i\tau_0 \log x) + O\{\exp(-c\sqrt{\log x})\} \\
&\text{if } \pi' \cong \pi \otimes \alpha^{i\tau_0} \text{ for some } \tau_0 \in \mathbb{R}^\times; \\
&= c_6 + O\{\exp(-c\sqrt{\log x})\} \\
&\text{if } \pi' \not\cong \pi \otimes \alpha^{it} \text{ for any } t \in \mathbb{R}.
\end{aligned}$$

Here Ei is the exponential integral, and $c, c_4, \dots, c_6$ are positive constants.

Recall that

$$\text{Ei}(i\tau_0 \log x) = \frac{x^{i\tau_0}}{i\tau_0 \log x} \left(\sum_{k=0}^n \frac{k!}{(i\tau_0 \log x)^k} \right) + O((\log x)^{-n-2}).$$

Our Corollary 1.5 is thus in a more precise form than Selberg's Conjecture 1.1.

The error terms in our theorem and corollaries are in a form that reflects very much our present knowledge of zero free regions for our Rankin-Selberg L -functions (see §4). The proofs of Corollaries 1.4 and 1.5 proceed along standard arguments, based on variations of Abel summation. We will thus not give these proofs here, but only point out that in the proof of Corollary 1.5, Hypothesis H is used to control sums over prime powers in the expression on the left side of (1.6). This way we can obtain a sum taken over primes as in (1.7).

We would like to express our heartfelt thanks to F. Shahidi for information on zero-free regions of automorphic L -functions, to the referee for detailed, in depth comments and suggestions, and to P. Sarnak for constructive remarks on an early draft of this paper.

2. Rankin-Selberg L -functions. We will use the Rankin-Selberg L -functions $L(s, \pi \times \tilde{\pi}')$ as developed by Jacquet, Piatetski-Shapiro, and Shalika [4], Shahidi [20], and Mœglin and Waldspurger [12], where π and π' are automorphic irreducible cuspidal representations of GL_m and $GL_{m'}$, respectively, over $\mathbb{Q}$ with unitary central characters. This L -function is given by local factors:

$$(2.1) \quad L(s, \pi \times \tilde{\pi}') = \prod_p L_p(s, \pi_p \times \tilde{\pi}'_p)$$

where

$$L_p(s, \pi_p \times \tilde{\pi}'_p) = \prod_{j=1}^m \prod_{k=1}^{m'} (1 - \alpha_\pi(p, j) \bar{\alpha}_{\pi'}(p, k) p^{-s})^{-1}.$$

The Archimedean local factor $L_\infty(s, \pi_\infty \times \tilde{\pi}'_\infty)$ is defined by

$$L_\infty(s, \pi_\infty \times \tilde{\pi}'_\infty) = \prod_{j=1}^m \prod_{k=1}^{m'} \Gamma_{\mathbb{R}}(s + \mu_{\pi \times \tilde{\pi}'}(j, k))$$

where the complex numbers $\mu_{\pi \times \tilde{\pi}'}(j, k)$ satisfy the trivial bound

$$(2.2) \quad \operatorname{Re} \mu_{\pi \times \tilde{\pi}'}(j, k) > -1.$$

Denote

$$\Phi(s, \pi \times \tilde{\pi}') = L_\infty(s, \pi_\infty \times \tilde{\pi}'_\infty) L(s, \pi \times \tilde{\pi}').$$

We will need the following properties of the L -functions $L(s, \pi \times \tilde{\pi}')$ and $\Phi(s, \pi \times \tilde{\pi}')$.

RS1. The Euler product for $L(s, \pi \times \tilde{\pi}')$ in (2.1) converges absolutely for $\operatorname{Re} s > 1$ (Jacquet and Shalika [5]).

RS2. The complete L -function $\Phi(s, \pi \times \tilde{\pi}')$ has an analytic continuation to the entire complex plane and satisfies a functional equation

$$\Phi(s, \pi \times \tilde{\pi}') = \varepsilon(s, \pi \times \tilde{\pi}') \Phi(1-s, \tilde{\pi} \times \pi'),$$

with

$$\varepsilon(s, \pi \times \tilde{\pi}') = \tau(\pi \times \tilde{\pi}') Q_{\pi \times \tilde{\pi}'}^{-s}$$

where $Q_{\pi \times \tilde{\pi}'} > 0$ and $\tau(\pi \times \tilde{\pi}') = \pm Q_{\pi \times \tilde{\pi}'}^{1/2}$ (Shahidi [20], [21], [22], and [23]).

RS3. Denote $\alpha(g) = |\det(g)|$. When $\pi' \not\cong \pi \otimes \alpha^{it}$ for any $t \in \mathbb{R}$, $\Phi(s, \pi \times \tilde{\pi}')$ is holomorphic. When $m = m'$ and $\pi' \cong \pi \otimes \alpha^{i\tau_0}$ for some $\tau_0 \in \mathbb{R}$, the only poles of $\Phi(s, \pi \times \pi')$ are simple poles at $s = i\tau_0$ and $1 + i\tau_0$ coming from $L(s, \pi \times \tilde{\pi}')$ (Jacquet and Shalika [5], [6], Moeglin and Waldspurger [12]). We will only consider the latter case in the proof of Theorem 1.3.

RS4. $\Phi(s, \pi \times \tilde{\pi}')$ is meromorphic of order one away from its poles, and bounded in vertical strips (Gelbart and Shahidi [2]).

RS5. $\Phi(s, \pi \times \tilde{\pi}')$ and $L(s, \pi \times \tilde{\pi}')$ are non-zero in $\text{Re } s \geq 1$. (Shahidi [20])

3. Estimation of logarithmic derivatives. Let $\mathbb{C}(m)$ be the complex plane with the discs

$$|s - 2n + \mu_{\pi \times \tilde{\pi}'}(j, k)| < \frac{1}{8m^2}, \quad n \leq 0, \quad 1 \leq j, k \leq m,$$

excluded. Here we give a remark about the structure of $\mathbb{C}(m)$. For $j, k = 1, \dots, m$, denote by $\beta(j, k)$ the fractional part of $\text{Re}(\mu_{\pi \times \tilde{\pi}'}(j, k))$. In addition we let $\beta(0, 0) = 0$ and $\beta(m+1, m+1) = 1$. Then all $\beta(j, k) \in [0, 1]$, and hence there exist $\beta(j_1, k_1), \beta(j_2, k_2)$ such that $\beta(j_2, k_2) - \beta(j_1, k_1) \geq 1/(3m^2)$ and there is no $\beta(j, k)$ lying between $\beta(j_1, k_1)$ and $\beta(j_2, k_2)$. It follows that the strip $S_0 = \{s : \beta(j_1, k_1) + 1/(8m^2) \leq \text{Re } s \leq \beta(j_2, k_2) - 1/(8m^2)\}$ is contained in $\mathbb{C}(m)$. Consequently, for all $n = 0, -1, -2, \dots$, the strips

$$(3.1) \quad S_n = \{s : n + \beta(j_1, k_1) + 1/(8m^2) \leq \text{Re } s \leq n + \beta(j_2, k_2) - 1/(8m^2)\}$$

are subsets of $\mathbb{C}(m)$. This structure of $\mathbb{C}(m)$ will be used later.

In Liu and Ye [10] and [11], we proved the following lemmas. It is believed that a much sharper result than Lemma 3.1 could be obtained using Selberg's explicit formula. This and generalization to a wider class of L -functions will be studied in a subsequent paper.

LEMMA 3.1. Assume $m = m'$ and $\pi' \cong \pi \otimes \alpha^{i\tau_0}$ for some nonzero $\tau_0 \in \mathbb{R}$. (a) Let $T > 2$. The number $N(T)$ of zeros of $L(s, \pi \times \tilde{\pi}')$ in the region $0 \leq \text{Re } s \leq 1, |\text{Im } s| \leq T$ satisfies

$$N(T+1) - N(T) \ll \log(Q_{\pi \times \tilde{\pi}'} T)$$

and

$$N(T) \ll T \log(Q_{\pi \times \tilde{\pi}'} T).$$

(b) For any $|T| > 2$, we have

$$\sum_{|T - \operatorname{Im} \rho| > 1} \frac{1}{(T - \operatorname{Im} \rho)^2} \ll \log(Q_{\pi \times \tilde{\pi}'} |T|).$$

(c) Let $s = \sigma + it$ with $-2 \leq \sigma \leq 2, |t| > 2$. If $s \in \mathbb{C}(m)$ is not a zero of $L(s, \pi \times \tilde{\pi}')$, then

$$\begin{aligned} & \frac{d}{ds} \log L(s, \pi \times \tilde{\pi}') \\ &= \sum_{|t - \operatorname{Im} \rho| \leq 1} \frac{1}{s - \rho} - \frac{1}{s - 1 - i\tau_0} - \frac{1}{s - i\tau_0} + O(\log(Q_{\pi \times \tilde{\pi}'} |t|)). \end{aligned}$$

(d) For $|T| > 2$, there exists τ with $T \leq \tau \leq T + 1$ such that when $-2 \leq \sigma \leq 2$

$$\frac{d}{ds} \log L(\sigma + i\tau, \pi \times \tilde{\pi}') \ll \log^2(Q_{\pi \times \tilde{\pi}'} |\tau|).$$

(e) For $|T| > 2$, there exists τ with $T \leq \tau \leq T + 1$ such that when $-2 \leq \sigma \leq 2$

$$\frac{d^2}{ds^2} \log L(\sigma + i\tau, \pi \times \tilde{\pi}') \ll \log^3(Q_{\pi \times \tilde{\pi}'} |\tau|).$$

LEMMA 3.2. Assume $m = m'$ and $\pi' \cong \pi \otimes \alpha^{i\tau_0}$ for some nonzero $\tau_0 \in \mathbb{R}$ as before. If s is in some strip S_n as in (3.1) with $n \leq -2$, then

$$\frac{d^2}{ds^2} \log L(s, \pi \times \tilde{\pi}') \ll_m 1.$$

4. Zero free regions. We need a zero free region for the Rankin-Selberg L -function $L(s, \pi \times \tilde{\pi}')$ which was proved by Moreno [13] and [14]. See also Gelbart, Lapid, and Sarnak [1], and Sarnak [18]. In order for later usage, we formulate the theorem for automorphic L -functions attached to cuspidal representations of GL_m over an algebraic number field F . Similar formulation can also be made to Moreno's zero free region near the possible pole. As in [13] and [14], the constant c' in (4.1) below can be made more precise in terms of the infinite types of the representations.

Denote by q_v the number of elements in the residue field of F_v at a non Archimedean place v of F . Let π and π' be any automorphic irreducible cuspidal representations of $GL_m(F_{\mathbb{A}})$ and $GL_{m'}(F_{\mathbb{A}})$, respectively. Then their Rankin-Selberg L -function is defined by

$$L(s, \pi \times \tilde{\pi}') = \prod_{v < \infty} L_v(s, \pi_v \times \tilde{\pi}'_v)$$

where

$$L_v(s, \pi_v \times \tilde{\pi}'_v) = \prod_{j=1}^m \prod_{k=1}^{m'} (1 - \alpha_\pi(v, j) \bar{\alpha}_{\pi'}(v, k) q_v^{-s})^{-1}$$

for $\text{Re } s > 1$ and by analytic continuation to $\mathbb{C}$. We define its Archimedean local factors in the standard way. This Rankin-Selberg L -function satisfies the same properties **RS1** through **RS5**, and the lemmas in §3 also hold.

THEOREM 4.1. *Let π (resp. π') be any automorphic irreducible cuspidal representation of $GL_m(F_{\mathbb{A}})$ (resp. $GL_{m'}(F_{\mathbb{A}})$). Assume that at least one of π and π' is self-contragredient: $\pi \cong \tilde{\pi}$ or $\pi' \cong \tilde{\pi}'$. Then there is an effectively computable constant c' such that there is no zero of $L(s, \pi \times \tilde{\pi}')$ in the region*

$$(4.1) \quad \sigma \geq 1 - \frac{c'}{\log(Q|t| + 1)}, \quad |t| \geq 1.$$

Here Q is the largest of $Q_{\pi \times \tilde{\pi}}$, $Q_{\pi \times \tilde{\pi}'} = Q_{\pi' \times \tilde{\pi}}$, and $Q_{\pi' \times \tilde{\pi}'}$.

5. Proof of Theorem 1.3. We now prove Theorem 1.3 when $\pi' \cong \pi \otimes \alpha^{i\tau_0}$ for some $\tau_0 \in \mathbb{R}$. The proof for case of π and π' being not twisted equivalent, in particular, when $m \neq m'$, is the same with all terms related to τ_0 removed. By **RS1**, we have for $\text{Re } s > 1$ that

$$\frac{d}{ds} \log L(s, \pi \times \tilde{\pi}') = - \sum_{n=1}^{\infty} \frac{\Lambda(n) a_\pi(n) \bar{a}_{\pi'}(n)}{n^s},$$

and therefore

$$K(s) := \frac{d^2}{ds^2} \log L(s, \pi \times \tilde{\pi}') = \sum_{n=1}^{\infty} \frac{(\log n) \Lambda(n) a_\pi(n) \bar{a}_{\pi'}(n)}{n^s}.$$

By **RS3** and **RS5**, $K(s)$ is holomorphic in $\text{Re } s > 1$. On $\text{Re } s = 1$, $L(s, \pi \times \tilde{\pi}')$ is nonzero (**RS5**) and has only a simple pole at $s = 1 + i\tau_0$. Thus

$$(5.1) \quad K(s) = \frac{1}{(s - 1 - i\tau_0)^2} + G(s)$$

has only a double pole in $\text{Re } s \geq 1$, where $G(s)$ is analytic for $\text{Re } s \geq 1$. On $\mathbb{C}$, $K(s)$ also has a double pole at each of the pole at $i\tau_0$, trivial zeros, and nontrivial zeros of $L(s, \pi \times \tilde{\pi}')$.

By Conjecture 1.2 and (1.1), we have

$$|a_\pi(p^k)| \leq m, \quad |a_{\pi'}(p^k)| \leq m'.$$

Therefore,

$$|(\log n) \Lambda(n) a_\pi(n) \bar{a}_{\pi'}(n)| \leq mm' \log^2 n,$$

and for $\sigma > 1$,

$$\sum_{n=1}^{\infty} \frac{|(\log n)\Lambda(n)a_{\pi}(n)\bar{a}_{\pi'}(n)|}{n^{\sigma}} \ll \frac{1}{(\sigma-1)^2}.$$

Let $T = \exp(\sqrt{\log x})$ and set $b = 1 + 1/\log x$. By Perron's summation formula (see e.g. Theorem 5.1 in [7]), we have

$$\begin{aligned} & \sum_{n \leq x} (\log n)\Lambda(n)a_{\pi}(n)\bar{a}_{\pi'}(n) \\ &= \frac{1}{2\pi i} \int_{b-iT}^{b+iT} K(s) \frac{x^s}{s} ds + O\left(\frac{x^b}{T(b-1)^2}\right) + O\left(\frac{x \log^3 x}{T}\right) \\ (5.2) \quad &= \frac{1}{2\pi i} \int_{b-iT}^{b+iT} K(s) \frac{x^s}{s} ds + O\left(\frac{x \log^3 x}{T}\right). \end{aligned}$$

Here we used the Ramanujan Conjecture 1.2 to control the size of error terms in (5.2).

Choose a with $-2 < a < -1$ such that the line $\operatorname{Re} s = a$ is contained in the strip $S_{-2} \subset \mathbb{C}(m)$; this is guaranteed by the structure of $\mathbb{C}(m)$. Let T be the τ such that Lemma 3.1(e) holds, by adding a constant d with $0 < d < 1$ to our $T = \exp(\sqrt{\log x})$ if necessary. Now we consider the contour

$$C_1 : \quad b \geq \sigma \geq a, \quad t = -T;$$

$$C_2 : \quad \sigma = a, \quad -T \leq t \leq T;$$

$$C_3 : \quad a \leq \sigma \leq b, \quad t = T.$$

Note that the two poles, some trivial zeros, and certain nontrivial zeros of $L(s, \pi \times \tilde{\pi}')$, as well as the pole at $s = 0$ are passed by the shifting of the contour. The trivial zeros can be determined by the functional equation in **RS2**: $s = -\mu_{\pi \times \tilde{\pi}'}(j, k)$ with $a < -1 - \operatorname{Re}(\mu_{\pi \times \tilde{\pi}'}(j, k)) < 0$. Here we used the facts that $\operatorname{Re}(\mu_{\pi \times \tilde{\pi}'}(j, k)) > -1$ and $-2 < a < -1$.

Then we have

$$(5.3) \quad \frac{1}{2\pi i} \int_{b-iT}^{b+iT} K(s) \frac{x^s}{s} ds$$

$$(5.4) \quad = \frac{1}{2\pi i} \left(\int_{C_1} + \int_{C_2} + \int_{C_3} \right) K(s) \frac{x^s}{s} ds$$

$$(5.5) \quad + \operatorname{Res}_{s=0} K(s) \frac{x^s}{s}$$

$$(5.6) \quad + \operatorname{Res}_{s=i\tau_0, 1+i\tau_0} K(s) \frac{x^s}{s}$$

$$(5.7) \quad + \sum_{a+1 < -\operatorname{Re}(\mu_{\pi \times \tilde{\pi}'}(j, k)) < 1} \operatorname{Res}_{s=-\mu_{\pi \times \tilde{\pi}'}(j, k)} K(s) \frac{x^s}{s}$$

$$(5.8) \quad + \sum_{|\operatorname{Im} \rho| \leq T} \operatorname{Res}_{s=\rho} K(s) \frac{x^s}{s}.$$

By Lemma 3.1(e), we get

$$(5.9) \quad \int_{C_1} \ll \int_a^b \log^3(Q_{\pi \times \tilde{\pi}'} T) \frac{x^\sigma}{T} d\sigma \ll \frac{x \log^3(Q_{\pi \times \tilde{\pi}'} T)}{T \log x},$$

and the same upper bound also holds for the integral on C_3 . By Lemma 3.2, then

$$(5.10) \quad \int_{C_2} \ll \int_{-T}^T \frac{x^a}{|t|+1} dt \ll \frac{\log T}{x}.$$

Obviously, (5.5) is

$$(5.11) \quad \operatorname{Res}_{s=0} K(s) \frac{x^s}{s} = K(1).$$

Since the poles at $s = i\tau_0$ and $s = 1 + i\tau_0$ are double poles, the residues in (5.6) give

$$\lim_{s \rightarrow i\tau_0} \frac{d}{ds} (s - i\tau_0)^2 K(s) \frac{x^s}{s} + \lim_{s \rightarrow 1+i\tau_0} \frac{d}{ds} (s - 1 - i\tau_0)^2 K(s) \frac{x^s}{s}.$$

For the second term, we have, by (5.1),

$$(5.12) \quad \begin{aligned} & \lim_{s \rightarrow 1+i\tau_0} \frac{d}{ds} (s - 1 - i\tau_0)^2 K(s) \frac{x^s}{s} \\ &= \lim_{s \rightarrow 1+i\tau_0} \frac{d}{ds} (1 + (s - 1 - i\tau_0)^2 G(s)) \frac{x^s}{s} \\ &= \lim_{s \rightarrow 1+i\tau_0} \frac{d}{ds} \frac{x^s}{s} \\ &= \frac{x^{1+i\tau_0}}{1+i\tau_0} \log x - \frac{x^{1+i\tau_0}}{(1+i\tau_0)^2}. \end{aligned}$$

The first term can be estimated similarly, and we get

$$\lim_{s \rightarrow i\tau_0} \frac{d}{ds} (s - i\tau_0)^2 K(s) \frac{x^s}{s} = \lim_{s \rightarrow i\tau_0} \frac{d}{ds} \frac{x^s}{s} \ll \log x.$$

Consequently (5.6) is

$$(5.13) \quad \frac{x^{1+i\tau_0}}{1+i\tau_0} \log x - \frac{x^{1+i\tau_0}}{(1+i\tau_0)^2} + O(\log x).$$

Near a trivial zero $s = -\mu_{\pi \times \tilde{\pi}'}(j, k)$ of order l in (5.7), we can express $K(s)$ as $-l/(s + \mu_{\pi \times \tilde{\pi}'}(j, k))^2$ plus an analytic function, like in (5.1). The residues in (5.7) can therefore be computed similar to what we did in (5.12). By (2.2), we know that $\operatorname{Re}(\mu_{\pi \times \tilde{\pi}'}(j, k)) \geq \delta - 1$ for some $\delta > 0$. Consequently (5.7) is bounded

$$(5.14) \quad \sum_{a < -1 - \operatorname{Re}(\mu_{\pi \times \tilde{\pi}'}(j, k)) < 0} \operatorname{Res}_{s = -\mu_{\pi \times \tilde{\pi}'}(j, k)} K(s) \frac{x^s}{s} \ll x^{1-\delta} \log x.$$

To compute the residues corresponding to nontrivial zeros in (5.8), we note that $\Phi(s, \pi \times \tilde{\pi}')$ is of order 1 (**RS4**), and $\Phi(1, \pi \times \tilde{\pi}') \neq 0$ (**RS5**). Using a standard argument, we see that

$$\sum_{|\gamma| \leq T} \frac{1}{|\rho|} \ll \log^2 T.$$

Consequently, (5.8) becomes

$$\begin{aligned} \sum_{|\operatorname{Im} \rho| \leq T} \operatorname{Res}_{s=\rho} K(s) \frac{x^s}{s} &= - \sum_{|\operatorname{Im} \rho| \leq T} \operatorname{Res}_{s=\rho} \frac{1}{(s-\rho)^2} \frac{x^s}{s} \\ &\ll \sum_{|\operatorname{Im} \rho| \leq T} \left| \frac{x^\rho \log x}{\rho} \right|. \end{aligned}$$

Using Moreno's zero free region in Theorem 4.1, we get

$$(5.15) \quad \ll x \exp \left(-c' \frac{\log x}{\log T} \right) (\log x) \log^2 T$$

By taking $T = \exp(\sqrt{\log x}) + d$ for some d with $0 < d < 1$, we can bound (5.9), (5.10), (5.15), and the error term in (5.2) by $O\{x \exp(-\frac{c'}{2} \sqrt{\log x})\}$ for the c' in Theorem 4.1. Using the main term from (5.13) and other error terms from (5.10), (5.11), (5.14), and (5.15), we get a proof of Theorem 1.3. $\square$

REFERENCES

- [1] S.S. Gelbart, E.M. Lapid, and P. Sarnak, *A new method for lower bounds of L-functions*, C.R. Acad. Sci. Paris, Ser. I **339** (2004), 91-94.

- [2] S.S. Gelbart and F. Shahidi, *Boundedness of automorphic L -functions in vertical strips*, J. Amer. Math. Soc., **14** (2001), 79-107.
- [3] R. Godement and H. Jacquet, *Zeta functions of simple algebras*, Lecture Notes in Math., **260**, Springer-Verlag, Berlin, 1972.
- [4] H. Jacquet, I.I. Piatetski-Shapiro, and J. Shalika, *Rankin-Selberg convolutions*, Amer. J. Math., **105** (1983), 367-464.
- [5] H. Jacquet and J.A. Shalika, *On Euler products and the classification of automorphic representations I*, Amer. J. Math., **103** (1981), 499-558.
- [6] H. Jacquet and J.A. Shalika, *On Euler products and the classification of automorphic representations II*, Amer. J. Math., **103** (1981), 777-815.
- [7] A.A. Karatsuba, *Basic Analytic Number Theory*, Springer, Berlin 1993.
- [8] H. Kim, *Functoriality for the exterior square of GL_4 and the symmetric fourth of GL_2* , J. Amer. Math. Soc., **16** (2003), 139-183.
- [9] H. Kim and P. Sarnak, *Refined estimates towards the Ramanujan and Selberg conjectures*, Appendix to Kim [8].
- [10] Jianya Liu and Yangbo Ye, *Superposition of zeros of distinct L -functions*, Forum Math., **14** (2002), 419-455.
- [11] Jianya Liu and Yangbo Ye, *Weighted Selberg orthogonality and uniqueness of factorization of automorphic L -functions*, preprint.
- [12] C. Moeglin and J.-L. Waldspurger, *Le spectre résiduel de $GL(n)$* , Ann. Sci. École Norm. Sup., (4) **22** (1989), 605-674.
- [13] C.J. Moreno, *Explicit formulas in the theory of automorphic forms*, Lecture Notes Math. vol. **626**, Springer, Berlin, 1977, 73-216.
- [14] C.J. Moreno, *Analytic proof of the strong multiplicity one theorem*, Amer. J. Math., **107** (1985), 163-206.
- [15] M. Ram Murty, *Selberg's conjectures and Artin L -functions*, Bull. Amer. Math. Soc., **31** (1994), 1-14.
- [16] M. Ram Murty, *Selberg's conjectures and Artin L -functions II*, Current trends in mathematics and physics, Narosa, New Delhi, 1995, 154-168.
- [17] Z. Rudnick and P. Sarnak, *Zeros of principal L -functions and random matrix theory*, Duke Math. J., **81** (1996), 269-322.
- [18] P. Sarnak, *Non-vanishing of L -functions on $\Re(s) = 1$* , Shalika's 60th Birthday, preprint, 2003.

- [19] A. Selberg, *Old and new conjectures and results about a class of Dirichlet series*, Collected Papers, vol. II, Springer, 1991, 47-63.
- [20] F. Shahidi, *On certain L -functions*, Amer. J. Math., **103** (1981), 297-355.
- [21] F. Shahidi, *Fourier transforms of intertwining operators and Plancherel measures for $GL(n)$* , Amer. J. Math., **106** (1984), 67-111.
- [22] F. Shahidi, *Local coefficients as Artin factors for real groups*, Duke Math. J., **52** (1985), 973-1007.
- [23] F. Shahidi, *A proof of Langlands' conjecture on Plancherel measures; Complementary series for p -adic groups*, Ann. Math., **132** (1990), 273-330.

JIANYA LIU: DEPARTMENT OF MATHEMATICS, SHANDONG UNIVERSITY

JINAN 250100, CHINA.

jyliu@sdu.edu.cn

YANGBO YE: DEPARTMENT OF MATHEMATICS, THE UNIVERSITY OF IOWA

IOWA CITY, IOWA 52242-1419, USA.

yey@math.uiowa.edu