

## 12. Topological Spaces

Defn: A **topology** on a set  $X$  is a collection  $\mathcal{T}$  of subsets of  $X$  having the following properties:

- a.)  $\emptyset, X \in \mathcal{T}$ .
- b.)  $U_\alpha \in \mathcal{T}$  implies  $\cup U_\alpha \in \mathcal{T}$
- c.)  $U_i \in \mathcal{T}$  implies  $\cap_{i=1}^n U_i \in \mathcal{T}$

Defn:  $U$  is open if  $U \in \mathcal{T}$

Ex 2a: The discrete topology on  $X = \mathcal{P}(X) =$  set of all subsets of  $X$ .

Ex 2b: The indiscrete or trivial topology on  $X = \{\emptyset, X\}$ .

Ex 3: The finite complement topology on  $X = \mathcal{T}_f = \{U \mid X - U \text{ is finite or } X - U = X\}$ .

Ex 4: The countable complement topology on  $X = \mathcal{T}_c = \{U \mid X - U \text{ is countable or } X - U = X\}$ .

Defn: Suppose the  $\mathcal{T}$  and  $\mathcal{T}'$  are two topologies on  $X$  such that  $\mathcal{T} \subset \mathcal{T}'$ . Then  $\mathcal{T}'$  is **finer** or **larger** than  $\mathcal{T}$  and  $\mathcal{T}$  is **coarser** or **smaller** than  $\mathcal{T}'$ . If  $\mathcal{T}'$  properly contains  $\mathcal{T}$ , then  $\mathcal{T}'$  is **strictly finer** than  $\mathcal{T}$  and  $\mathcal{T}$  is **strictly coarser** than  $\mathcal{T}'$ .

Defn:  $\mathcal{T}$  is **comparable** with  $\mathcal{T}'$  if either  $\mathcal{T} \subset \mathcal{T}'$  or  $\mathcal{T}' \subset \mathcal{T}$ .

---

## 13: Basis for a Topology

Defn: If  $X$  is a set, a **basis** for a topology on  $X$  is a collection  $\mathcal{B}$  of subsets of  $X$  (called **basis elements**) such that

- (1) For each  $x \in X$ , there is at least one basis element  $B$  containing  $x$ .
- (2) If  $x \in B_1 \cap B_2$  where  $B_1, B_2 \in \mathcal{B}$ , then there exists  $B_3 \in \mathcal{B}$  such that  $x \in B_3 \subset B_1 \cap B_2$ .

The topology  $\mathcal{T}$  generated by a basis  $\mathcal{B}$  is defined as follows:  $U$  is open if and only if for all  $x \in U$ , there exists  $B \in \mathcal{B}$  such that  $x \in B \subset U$

Example 1a: The set of all open intervals in  $R$  is a basis for a topology on  $R$  (the standard topology).

Example 1b: The set of all open circular regions in  $R^2$  is a basis for a topology on  $R^2$  (the standard topology).

Example 2: The set of all open rectangular regions in  $R^2$  is a basis for a topology on  $R^2$  (the standard topology).

Note the basis in Example 1b and the basis in Example 2 both generated the same topology.

Example 3:  $\{x \mid x \in X\}$  is a basis for the discrete topology on  $X$ .

Lemma 13.1: Let  $\mathcal{B}$  be a basis for a topology  $\mathcal{T}$  on  $X$ . Then  $\mathcal{T}$  = set of all unions of elements of  $\mathcal{B}$ .

Lemma 13.2: Let  $X$  be a topological space. Suppose that  $\mathcal{C}$  is a collection of open sets of  $X$  such that for each open set  $U$  of  $X$  and each  $x \in U$ , there is an element  $C \in \mathcal{C}$  such that  $x \in C \subset U$ . Then  $\mathcal{C}$  is a basis for the topology on  $X$ .

Lemma 13.3: Let  $\mathcal{B}$  and  $\mathcal{B}'$  be a basis for  $\mathcal{T}$  and  $\mathcal{T}'$ , respectively, on  $X$ . Then the following are equivalent:

- (1)  $\mathcal{T}'$  is finer than  $\mathcal{T}$ .
- (2) For each  $x \in X$  and each basis element  $B \in \mathcal{B}$  containing  $x$ , there is a basis element  $B' \in \mathcal{B}'$  such that  $x \in B' \subset B$ .

Defn:

1.)  $\mathcal{B} = \{(a, b) \mid a, b \in R, a < b\}$  is a basis for the standard topology on  $R$ .

2.)  $\mathcal{B}' = \{[a, b) \mid a, b \in R, a < b\}$  is a basis for the lower limit topology on  $R$ . When  $R$  has this topology, we denote it by  $R_l$ .

3.) Let  $K = \{\frac{1}{n} \mid n \in Z_+\}$ .  $\mathcal{B}'' = \mathcal{B} \cup \{(a, b) - K \mid a, b \in R, a < b\}$  is a basis for the K-topology on  $R$ . When  $R$  has this topology, we denote it by  $R_K$ .

Lemma 13.4: The topologies  $R_l$  and  $R_K$  are strictly finer than the standard topology, but they are not comparable with one another. ■

Definition: A **subbasis**  $\mathcal{S}$  for a topology on  $X$  is a collection of subsets of  $X$  whose union equals  $X$ . The **topology generated by the subbasis**  $\mathcal{S}$  is defined to be the collection  $\mathcal{T}$  of all unions of finite intersections of elements of  $\mathcal{S}$ .

Lemma: If  $\mathcal{S}$  is a subbasis for a topology on  $X$ , then  $\mathcal{B} =$  the set of all finite intersections of elements of  $\mathcal{S}$  is a basis for this topology.

HW p83: 4, 8

---

#### 14: The Order topology

(p. 24) A relation  $<$  on a set  $A$  is called an **order relation** (or a **simple order** or **linear order**) if it has the following properties:

- (1) (Comparability) For every  $x, y \in A$  for which  $x \neq y$ , either  $x < y$  or  $y < x$ .
- (2) (Nonreflexivity) For no  $x \in A$  does the relation  $x < x$  hold.
- (3) (Transitivity) If  $x < y$  and  $y < z$ , then  $x < z$ .

Defn: Let  $X$  be a set with a simple order relation. Assume that  $X$  has more than one element. Let  $\mathcal{B}$  be the collection of all sets of the following types:

- (1) All open intervals  $(a, b)$  in  $X$ .
- (2) All intervals of the form  $[a_0, b)$ , where  $a_0$  is the smallest element (if any) of  $X$ .
- (3) All intervals of the form  $(a, b_0]$ , where  $b_0$  is the largest element (if any) of  $X$ .

The collection  $\mathcal{B}$  is a basis for a topology on  $X$  which is called the **order topology**.

Note: If  $X$  has no smallest element, there are no sets of type (2). If  $X$  has no largest element, there are no sets of type (3).

Ex. 0: The order topology on  $(0, 1) \cup \{5\}$

Ex. 1: The order topology on  $R$  is the standard topology on  $R$ .

Ex. 2:  $R \times R$  in the dictionary order.

Ex. 3: Order topology on  $Z_+ =$  discrete topology.

Ex. 4: The order topology on  $X = \{1, 2\} \times Z_+$  is NOT the discrete topology.