

22M:132: Topology Final Exam

Dec. 17, 2008

1.) Let \mathbf{R} be the set of real numbers and let \mathbf{Z} be the set of integers. Let $\bar{d}(x, y) = \min\{1, |x - y|\}$, Identify the following subsets of \mathbf{R} .

[3] 1a.) $B_{\bar{d}}(0, 1) =$ _____

[3] 1b.) $\overline{B_{\bar{d}}(0, 1)} =$ _____

[3] 1c.) $\{x \mid \bar{d}(x, 0) \leq 1\} =$ _____

[3] 1d.) The set of limit points of $\mathbf{Z} = \mathbf{Z}' =$ _____

[3] 1e.) The closure of $\mathbf{Z} = \bar{\mathbf{Z}} =$ _____

[2] 2.) An example of a paracompact space is _____

3.) Circle T for true and F for false.

[2] 3a.) If X is paracompact, then an arbitrary union of closed sets is closed. T F

[2] 3b.) If \mathcal{A} is a locally finite collection of closed subsets of X , then $\cup_{A \in \mathcal{A}} A$ is closed. T F

[80] Prove 4 of the following 6. Clearly indicate your choices. Note \mathbf{R} is the set of real numbers

Your 4 choices: _____

1.) Let X be a topological space in which one-point sets are closed in X . Show that X is regular if and only if for all $x \in X$, for every open set U in X such that $x \in U$, there is an open set V such that $x \in V \subset \overline{V} \subset U$.

2i.) Suppose $f : X \rightarrow Y$ is bijective and continuous, X is compact, and Y is T_2 . Show that f is a homeomorphism.

ii.) Give an example of a function $f : X \rightarrow Y$ which is bijective and continuous, but not a homeomorphism where X is a subspace of a manifold and Y is a compact manifold.

3.) A connected, locally pathwise connected space is pathwise connected. (Hint: find a set which is both open and closed).

4.) Recall that if G is a topological group, then $m : G \times G \rightarrow G$, $m(x, y) = xy$ is continuous. Let G be a topological group and let $x, y \in G$.

i.) Show that for every open neighborhood U of xy , there exists open sets, V and W , such that $x \in V$, $y \in W$ and $VW \subset U$.

ii.) If U is an open set containing the identity element e , then there exists an open set V such that $e \in V$ and $V^2 = \{v_1v_2 \mid v_i \in V\} \subset U$.

5.) Every closed subspace of a paracompact space is paracompact.

6.) Let $Y^X = \{f : X \rightarrow Y\}$. Let $S(x, U) = \{f \in Y^X \mid f(x) \in U\}$.

The topology of pointwise convergence on Y^X is the topology generated by the subbasis $\mathcal{S} = \{S(x, U) \mid x \in X, U \text{ open in } Y\}$.

i.) $S(0, (1, 2) \times (1, 2)) \subset (\mathbf{R}^2)^{\mathbf{R}}$. Give an example of a function in $S(0, (1, 2) \times (1, 2))$

ii.) Prove that the sequence f_n converges in Y^X where Y^X has the topology of pointwise convergence if and only if for all $x \in X$, the sequence $f_n(x)$ converges to $f(x)$ in Y .